

Product Name:		Territory:		Colours:	1. Draft	23.09.2020, 15:30	Mock up
Vaminolact		MY		• Schwarz • Die cut	1. Corr	24.09.2020, 09:15	
Type of Packaging:	Dosage:				2. Corr		
Leaflet	100ml/500ml				3. Corr		
Material number:	2-D-Matrix Code:						
M0xxxx/00-MY	M0xxxx/00-MY						
Pharma-Code (Laetus):	EAN-Code:						
-	-						
Dimension:	Font:	Size:	Operator:				
180 x 294 mm	Antique Olive	9	Roberto Grill				

Vaminolact™

Description

Amino acid solution for intravenous nutrition.

Vaminolact™ is an electrolyte-free, clear, colourless to slightly yellow solution of amino acids, for intravenous nutrition in infants and children.

Composition

Amino acids (g/l):

Alanine	6.3
Arginine	4.1
Aspartic acid	4.1
Cysteine/cystine	1.0
Glutamic acid	7.1
Glycine (amino acetic acid)	2.1
Histidine	2.1
Isoleucine	3.1
Leucine	7.0
Lysine	5.6
Methionine	1.3
Phenylalanine	2.7
Proline	5.6
Serine	3.8
Taurine	0.3
Threonine	3.6
Tryptophan	1.4
Tyrosine	0.5
Valine	3.6

Water for injections to 1000 ml.

Total amount of amino acids: 65.3 g/l of which 31.9 g, including cysteine, tyrosine and histidine, are essential.

Nitrogen content: 9.3 g/l

Electrolytes: None

Antioxidant additives: none

Energy content: 1.0 MJ (240 kcal)

pH: 5.2

Osmolality: 510 mosmol per kg water

Pharmacodynamics

The amino acid composition of VAMINOLACT is based on that of human milk. Apart from their nutritive properties, the amino acids should have no specific pharmacodynamic effects when administered at recommended rates.

To optimize the utilization of supplied amino acids, adequate energy should be supplied in the form of carbohydrates (preferable glucose) and fat.

Pharmacokinetics

The pharmacokinetic properties of the infused amino acids are essentially the same as for amino acids supplied by ordinary food. However, the amino acids of dietary protein first enter the portal vein and then the systemic circulation, while intravenously infused amino acids reach the systemic circulation directly.

Indications

Vaminolact is indicated as a source of amino acids in paediatric patients needing intravenous nutrition.

Contraindications

Vaminolact is contra-indicated in patients with inborn errors of amino acid metabolism, irreversible liver damage and in severe uremia when dialysis facilities are not available.

Caution

Intravenous infusion of amino acids may be accompanied by increased urinary excretion of the trace elements copper and, in particular zinc, which should be taken into account in the dosing of trace elements, particularly during long-term intravenous nutrition.

Light exposure of solutions for intravenous parenteral nutrition, especially after admixture with trace elements and/or vitamins, may have adverse effects on clinical outcome in neonates, due to generation of peroxides and other degradation products. When used in neonates and children below 2 years, Vaminolact should be protected from ambient light until administration is completed (see sections Dosage and Administration).

Adverse events

Nausea occurs rarely. Transient increases in liver functions tests during intravenous nutrition have been reported. The reasons are at present unclear. The underlying disease, the components and their amounts in the intravenous feeding regimens have been suggested.

Hypersensitivity reactions have been reported with amino acid solutions. As with all hypertonic infusion solutions, thrombophlebitis may occur when peripheral veins are used. The incidence may be reduced by the simultaneous infusion of Intralipid.

Dosage and administration

Recommended dosage range:

Preterm neonates: 38 to 54 mL/kg/d (corresponding to 2.5 to 3.5 g amino acids/kg/d).

Gradual dose increase during the first days of infusion should be used, starting with a dose such as 23 to 38 mL/kg/d (corresponding to 1.5 to 2.5 g amino acids/kg/d) on the first day and increasing to 38 to 54 (corresponding to 2.5 to 3.5 g amino acids/kg/d) on the second day onwards.

Term neonates: 23 to 46 mL/kg/d (corresponding to 1.5 to 3.0 g amino acids/kg/d).

Gradual dose increase to the target dose should be used during the first days of infusion.

Infants: 15 to 38 mL/kg/d (corresponding to 1.0 to 2.5 g amino acids/kg/d).

Children and adolescents: 15 to 31 mL/kg/d (corresponding to 1.0 to 2.0 g amino acids/kg/d).

The duration of infusion should be at least 8 hours, preferably 12 hours as cyclic infusion or 24 hours as continuous infusion. In neonates and infants, the recommended duration of continuous infusion is 24 hours/d.

Vaminolact may be infused into the same central or peripheral vein as glucose and/or fat emulsion.

When used in neonates and children below 2 years, the solution (in bags and administration sets) should be protected from light exposure until administration is completed (see section Caution).



Interactions with Other Medications

Not to be expected.

Pregnancy and Lactation

Not applicable

Overdosage

Vomiting, flushing and sweating have been observed if VAMINOLACT is administered at a higher rate than recommended. In case of symptoms due to overdose, the infusion should be slowed down or discontinued. In case of symptoms due to overdose, the infusion should be slowed down or discontinued.

Compatibility

Vaminolact can be mixed with other nutrition solutions (lipid emulsions, carbohydrates, electrolyte solutions, trace elements and vitamins). Compatibility data can be obtained upon request.

Adding of solutions should be made aseptically.

Storage

Do not store above 30 °C. Do not freeze.

Instruction for Use

Vaminolact should be used immediately after opening, using sterile infusion devices. Any unused solution should be discarded.

Packsizes

Bottles of 100 ml and 500 ml.

Date of revision: April 2021

Manufactured by
Fresenius Kabi Austria GmbH, Graz, Austria

