

(L)148.5mm

(W)210mm

pharmaniaga™

Aneurine

 Tablet

CONTRAINDICATIONS

Risk benefit should be considered when the following problems exist.

- Sensitivity to thiamine
- Wernicke's encephalopathy

ADVERSE REACTIONS

Rare, especially in person with normal renal function. However, skin rash or itching may occur in individuals who are allergic to thiamine.

PRECAUTIONS

Alcohol may inhibit absorption of thiamine.

PREGNANCY AND LACTATION

This product is not intended for use in pregnant or lactating women.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

No studies on the effect on the ability to drive and use machine have been performed.

However, patients should be cautioned to see how they react before driving or operating machinery.

DOSAGE AND ADMINISTRATION

Adult:

For mild chronic deficiency – 5 mg to 30 mg daily. In severe thiamine deficiency doses of up to 300 mg daily have been given.

ROUTE OF ADMINISTRATION

Oral.

OVERDOSAGE

Overdosage with Thiamine is quite unlikely when administered by mouth.

STORAGE CONDITIONS

Store below 30°C. Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

Aneurine Tablet 3 mg

In blisters of 6930 tablets.

Aneurine Tablet 10 mg

In blisters of 490 and 6930 tablets.

Date of revision: 3rd September 2018

DESCRIPTION

Aneurine Tablet 3 mg

Light yellow, round, flat beveled edge tablet with 'KK' marking on one side and plain on the other.

Aneurine Tablet 10 mg

Light pink, round, flat beveled edge tablet with 'KK' marking on one side and plain on the other.

COMPOSITION

Each tablet contains Thiamine Mononitrate 3 mg or 10 mg.

ACTIONS

Thiamine is necessary for carbohydrate metabolism. It combines with the pyrophosphoric acid in nucleated cells, particularly in the liver, kidneys and white blood cells to form thiamine pyrophosphate (cocarboxylase), which acts as a coenzyme in such reactions as the decarboxylation of alpha-ketoacids, particularly of pyruvate and alpha-ketoglutarate. In the presence of thiamine deficiency, pyruvic and lactic acids accumulate in the tissues.

Thiamine pyrophosphate also acts as a coenzyme in the direct oxidative pathway of glucose metabolisms.

Thiamine is readily absorbed from the gastrointestinal tract, except in malabsorption syndromes. Thiamine is absorbed mainly in the duodenum. Alcohol inhibits absorption of thiamine. It is excreted as unchanged drug and metabolites in the urine.

INDICATIONS

Prevention and treatment of thiamine deficiency state, which may occur as a result of inadequate nutrition.

PRODUCT REGISTRATION HOLDER/ MANUFACTURER:

PHARMANIAGA MANUFACTURING BERHAD (60016-D)

11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

XXXXXXXXXXXX

attn

customer Pharmaniaga Manufacturing Berhad

date 22.03.18

1 colours

process colours:

Black

Please Chop & Sign For Approval

size (L) 148.5 x (W) 210mm

description Aneurine Leaflet



FocusPrint SDN BH D

t: 603-8766 6030 f: 603-8766 6033 (Factory)
website: www.focusprint.info

artwork prepared by: cynthia yap email: graphic@focusprint.info (Graphic Dept)

date:

IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.

ARTWORK LOG

Version no.	Date	Reason for Change
01	26.07.17	- Amend wording and add in Date of revision.
02	02.08.17	- Rearrange wording.
03	22.03.18	- Amend wording.