

Lampiran L1E
SISIP BUNGKUSAN PINDAAN ZHENG GU SHUI





正骨水

只供外用

成份 每100毫升此产品含以下药材提取物:

五马巡城	1.170毫克	鸡骨香	1.425毫克
白芷	1.235毫克	千斤拔	1.140毫克
大力王	1.140毫克	田七	2.375毫克
薄荷脑	0.093毫克	樟脑	0.190毫克

主治 传统用于解除肿胀及舒缓扭伤。

用法 用药棉蘸药液，先擦患处周围皮肤，作局部止痛后，然后用药棉渗透敷贴于，用纱布轻扎，上肢骨骼敷药一小时，下肢骨骼敷药二小时，即可将药棉去掉，以后每日擦药二至三次，直至痊愈。

指示 1. 不可内服。
2. 儿童皮肤柔软嫩，敷药时应将年龄配量减短，成年人亦应按用法掌握，以免药力过度刺激，引起皮肤发炎。
3. 如患处皮肤破血出血先止血，然后用此药擦于伤口周围，忌擦伤口。
4. 敷药后不可在患处包扎过紧，以免循环障碍引起水泡。

注意 1. 如病情无好转，请向医生咨询。
2. 此乃传统药。
3. 请将药物远离小孩。
4. 置于阴凉干燥处。
5. 此产品只供外用。
6. 本品挥发性强，用后应密封，不宜靠近高温及火处。
7. 可能会引起抽感。
8. 二岁以下婴儿不可使用。
9. 年龄较大的儿童使用时应谨慎。
10. 勿直接用于鼻孔处。
11. 勿接触眼睛。
12. 勿直接擦于伤口。



Imported by Product Registration Holder:
星马提煉药(马)有限公司 (Co. No. 11394 U)
SINMA MEDICAL PRODUCTS (M) SDN BHD.
No. 13, Jalan Teras 1,
Kawasan Industri Teras Bahang,
43300 Bahang, Selangor D.E., Malaysia.







ZHENG GU SHUI

(Liquid)

For External Use Only

Ingredients: Every 100ml of this product contains extraction of active ingredients as shown below:

Rhizoma Croton Tiglium	1.710mg
Radix Cinnamomum Camphora Nees et Ebern.	1.425mg
Radix Angelicae Dahuricae	1.235mg
Radix Moganiana Macrophylla	1.140mg
Rhizoma Inula Cappa	1.140mg
Radix Pseudo Ginseng	2.375mg
Menthol	0.093mg
Camphor	0.190mg

Indication: Traditionally used for reducing swelling and relieving sprains.

Directions: Rub the area peripheral to the wound with cotton soaked with the tincture for analgesia. Dress with soaked cotton and bandage gently. After one hour of medication on upper limbs or one and a half hours on lower limbs, remove the cotton. Apply two or three times daily thereafter until total recovery.

Precautions: 1. Not to be taken internally.
2. For children, duration of application should be proportionally reduced, in order to avoid over-stimulation of skin and inflammation.
For adults, also follow the above direction accordingly.
3. In case of open wound, bleeding must be stopped first. Then rub with this medicine around the wound. Never rub into the wound.
4. Never bandage the wounded area tightly, in order to prevent disturbance of circulation and blister-formation.

Attention: 1. If symptoms persist, please consult a doctor.
2. This is a traditional medicine.
3. Keep out of reach of children.
4. Store below 30°C and protect from light and moisture.
5. For external use only.
6. This medicine is highly volatile. The bottle should be tightly closed after use and keep away from fire and high temperature.

Warning: 1. Can cause convulsion.
2. Contraindicated in infants below 2 years of age.
3. Caution must be exercised when older children are treated.
4. Avoid direct application into nostrils.
5. Avoid contact with the eyes.
6. Do not apply to wounds or damaged skin.

It is dangerous to place any camphor-containing product into the nostrils of children. A small amount applied this way may cause immediate collapse.

Packing size: Available in 30ml, 100ml & 60ml (spray).

Shelf life: 5 years

Jauhi dari kanak-kanak **MAL19962254T**

Manufactured by:
广西玉林制药有限公司
GUANGXI YULIN PHARMACEUTICAL FACTORY
Cheng Zhan Road, Yulin, Guangxi, China.

