

ASUMALIFE[®] SYRUP

Ketotifen exerts potent anti-anaphylactic and antihistaminic effects. Its powerful and sustained H₁-receptor-blocking activity can be clearly dissociated from its anti-anaphylactic properties. Ketotifen has been used in the prophylactic treatment of asthma.

Description(s):

A clear, colorless solution with pineapple flavor.

Ingredient(s):

Each 5ml contains:

Ketotifen Fumarate 1.38mg
(eq. to Ketotifen 1mg)

Preservative:

Sodium Benzoate 1.875mg

Pharmacodynamics :

1. The anti-anaphylactic property of Ketotifen is due to its inhibition of the release of such chemical mediators as histamine and, in particular, SRS-A (the slow reacting substance of anaphylaxis which is a mixture of leukotrienes) from tissue mast cells, basophils, and pulmonary tissue; inhibition of bronchoconstriction due to SRS-A and inhibition of cellular Calcium ion uptake.
2. Ketotifen has been shown to reverse experimentally-induced tachyphylaxis to Isoprenaline.

Pharmacokinetic(s):

1. Ketotifen is absorbed from the gastrointestinal tract and is excreted in the urine and feces. It is eliminated biphasically with a short half-life of 3 - 5 hours and a longer one of 21 hours. Its main metabolite is the inactive Ketotifen-N-glucuronide.
2. Maximum plasma concentrations are reached within 2 - 4 hours.

Indication(s):

Prophylactic treatment of bronchial asthma and symptomatic treatment of allergic rhinitis, dermatitis, urticaria and pruritus. It is not effective in aborting established attacks of asthma.

Dosage and Administration:

Children 6 months- 3 years of age: 0.25ml per kg body weight twice daily

Adult and children over 3 years of age: 5ml twice daily

Patients known to be easily sedated should begin treatment with 2.5-5ml at night for the first few days.

To be taken with food. To be dispensed on physician's prescription.

Route of administration: Oral

Contraindication(s):

1. Contraindicated in patients with known hypersensitivity to Ketotifen.
2. Ketotifen should not be used for the treatment of acute asthma attacks.
3. Ketotifen should not be given to premature infants or neonates because of the higher risk of adverse effects with Ketotifen.

Side Effect(s) / Adverse Reaction(s):

Sedation, dry mouth, dizziness, increased appetite and weight gain.

Drug Interaction(s):

It may potentiate the effects of sedatives, hypnotics, antihistamines and alcohol. Patients should be advised to avoid alcoholic drinks.

Precaution(s) / Warning(s):

1. Concomitant use of Ketotifen with oral antidiabetic agents should be avoided as a reversible fall in the platelets count has been observed.
2. The dose should be reduced gradually over a period of one week upon withdrawal to prevent the recurrence of the symptoms of asthma.

3. Care should be exercised when driving a vehicle or operating machinery because Ketotifen may cause sedation.
4. Ketotifen, like most drugs, should be given to pregnant women and nursing mothers only under compelling circumstances.
5. If intercurrent infections occur, treatment must be supplemented by specific antimicrobial therapy. Patients should be advised to avoid alcoholic drinks.

Symptoms and Treatment for Overdosage, and Antidote(s):

Symptoms: Drowsiness to severe sedation; confusion and disorientation; tachycardia and hypotension; convulsions, especially in children; hyperexcitability in children; reversible coma. In most cases of acute toxicity, gastric lavage was performed following which patients made full recovery with only supportive treatment.

Shelf-Life:

2 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Packing(s):

Plastic bottle of 60ml and 100ml.

Plastic bottle of 1L and 4L (for export only).

SIRAP ASUMALIFE®

Bahan Aktif:

Setiap 5ml mengandungi:

Ketotifen Fumarate 1.38mg
(setara dengan Ketotifen 1mg)

Pengawet:

Sodium Benzoate 1.875mg

Indikasi:

Rawatan profilaktik untuk asma bronkial dan rawatan gejala untuk rinitis alergik, dermatitis, urtikaria dan pruritus. Ubat ini tidak berkesan untuk menghilangkan serangan asma yang sudah terbentuk.

Dos:

Kanak-kanak berumur 6 bulan-3 tahun: 0.25ml per kg berat badan 2 kali sehari

Orang dewasa dan kanak-kanak berumur 3 tahun ke atas: 5ml 2 kali sehari.

Pesakit-pesakit yang didapati mudah mengantuk apabila diberi ubat ini, dos rawatan seharusnya mula dengan 2.5ml-5ml diberi pada waktu malam untuk beberapa hari.

Diberi bersama dengan makanan. Didispen mengikut preskripsi doktor.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679