

건 명	알부민 20% 50mL 인서트(말레이시아 등록용)		
규 격 (가로x세로)	100x267mm	Q A	오창
		표시문안	4,5pt
색 상	먹 1도	후 가 공	
업체제작일	2025.07.03.	작 업 처	(주)트리스톤



칼선 PANTONE 185C

※ 인쇄진행시, 인쇄사양과 다를경우
녹십자로 즉시 연락주시기 바랍니다.

Normal Serum Albumin, Human Albumin-GCC 20%

Description

ALBUMIN-GCC(20%) is a sterile aqueous solution containing 20% purified human serum albumin(weight/volume). ALBUMIN-GCC(20%) is prepared by the Cohn ethanol blood fractionation method from pooled human plasma with a poor salt content obtained from healthy blood donors.
The product is stabilized with Sodium caprylate and Acetyl tryptophan. ALBUMIN-GCC(20%) is osmotically equivalent to four times its volume of normal citrated plasma. A liter of ALBUMIN-GCC(20%) solution contains 113,5-138,7mM of sodium ion. The product contains no preservatives.
ALBUMIN-GCC(20%) is heated at 60°C for ten hours. This heat treatment completely destroys the causative agents of viral hepatitis(HBV, HCV) and HIV -1 and -2. This product is prepared from units of human plasma that have been tested and found nonreactive for HBsAg and antibodies to HIV -1 and -2 and HCV.

Therapeutic class : Blood Fractions

Composition

Each 50ml vial contains
Normal Human Serum Albumin(active ingredient) ----- 10g
Sodium caprylate(as stabilizer) ----- 0,13295g
Acetyl tryptophan(as stabilizer) ----- 0,197g
Distilled water for injection ----- q.s.

Action and Mode or Mechanism of Action

The primary function of Albumin is that of providing colloidal osmotic pressure within the circulation. The intravenous administration of ALBUMIN-GCC results, therefore, in drawing extravascular fluid into the vascular system. Serum albumin which contributes to the buffer capacity of the plasma, has transport function. Although Albumin contains some amino acids, it provides only moderate nutritive effect.

Indication

- 1) Hypoalbuminemia caused by albumin deficiency(Burn, Nephrotic Syndrome) and disfunction of albumin synthesis
- 2) Hemorrhagic shock

Recommended Dosage, Dosage Schedule and Route of Administration

ALBUMIN-GCC(20%) is administered intravenously.
Hypoalbuminemia caused by albumin deficiency(Burn, Nephrotic Syndrome) and disfunction of albumin synthesis.
Inject intravenously
The total dosage will vary with clinical condition and size of the individual
• In adults
An initial dose of 100ml infusion is suggested.
Additional amount may be administered as clinically indicated.
• In children
A dose one quarter to one half the adult dose or dosage may be calculated on the basis of 1,25-2,5ml per kilogram of body weight.
Hemorrhagic Shock
In the treatment of the patient in shock with greatly reduced blood volume, ALBUMIN-GCC may be administered intravenously as rapidly as necessary in order to improve the clinical condition and restore normal blood volume. This may be repeated in 15-30minutes if the initial dose fails to prove adequate. In the patient with a slightly low or normal blood volume, the rate of administration is not more than 1ml per minute. The usual rate of administration in children should be calculated on one quarter to one half the adult rate

Contraindication

ALBUMIN-GCC(20%) is contraindicated in patients with severe anemia or cardiac failure in the presence of normal or increased intravascular volume, and with known hypersensitivity to commercially available albumin solutions.

Adverse Reaction

Allergic or pyrogenic reactions are characterized primarily by fever and chills; rash, nausea, vomiting, tachycardia and hypotension have also been reported. ALBUMIN-GCC(20%), particularly if administered rapidly, may result in vascular overload with resultant pulmonary edema.

Precaution/Warning

ALBUMIN-GCC(20%) should be administered with caution to patients with low cardiac reserve.
Rapid infusion may cause vascular overload with resultant pulmonary edema.
Patients should be closely monitored for signs of increased venous pressure.
• Dehydrated patients may require additional fluids.
• Rapid administration to injured patients may result in vascular overload or bleeding from previously undetected sites.
• Use in pregnancy : There is very little experience of the use of albumin in the early stages of pregnancy.

Special Warning

Although this blood product is made from plasma which has been screened or tested for some known transmissible agents including HIV(AIDS), hepatitis B and hepatitis C, the limited sensitivity and reliability of such tests and screening procedures and possible contamination with other or unknown agents must be borne in mind.
Chemical processing and virus inactivation stages included in the manufacture of this product are believed to render it safe from the risk of infection. Nevertheless this possibility must always be considered, and should be conveyed, whenever possible, to patients who may receive the product.

Spillage or Breakage

Do not use if a break of the container or spillage occur. Precautions should be taken to avoid contamination of cuts and abrasions, as well as to avoid inhalation or swallowing of the spillage.
Adequate disinfection can be obtained with the application of 1% sodium hypochlorite solution for 15minutes. Commercial bleaches may be diluted appropriately to obtain this concentration. Please contact the supplier if you have any problem.

Symptoms and Treatment For Overdosage And Antidote(s)

Toxicity and treatment of overdosage : Administration of large quantities of albumin should be supplemented with red cell concentrates or replaced by whole blood to combat the relative anaemia which would follow such use.
If circulatory embarrassment or pulmonary edema should develop, the infusion must be stopped immediately and specific treatment given.

User Instruction:

- Cleanse exposed rubber stopper with a suitable germicidal solution and prepare the administration equipment as follows:
- a) Close clamp on infusion set.
 - b) With bottle upright, thrust piercing pin straight through stopper center. Do not twist or angle.
 - c) Immediately invert bottle to automatically establish proper fluid level in drip chamber(half full).
 - d) Attach infusion set, open clamp and solution to expel air from tubing and needle, then close clamp.
 - e) Make venipuncture and adjust flow.
 - f) Discard all administration equipment after use.

Pharmaceutical Precautions:

Should not be used if they appear turbid or if there is sediment in the bottle, or more than 4 hours after the container has been penetrated. Any remaining contents should be discarded.

Storage Conditions:

Store below 30°C. Do not freeze.

Expiry Date:

39months from the manufacturing date

Packing/Pack Size

50ml vial(20% Albumin)

Name and Address of Product Registration Holder in Malaysia.

PROPHARM(M) SDN. BHD.
8, Jln. Udang Harimau 2
Medan Niaga Kepong
51200 Kuala Lumpur
Tel : 03-62436389

Manufactured and Released by:



GC Biopharma

GC Biopharma Corp.
586, Gwahaksaneop 2-ro, Ochang-eup, Cheongwon-gu,
Cheongju-si, Chungcheongbuk-do, Korea

Ins-02-211-4
Revised on 09-01-2024