

安全藥油

安全藥油是一种常见的家用药物，实用方便，随身携带。这是一种特殊的制剂，可以减轻疼痛和酸痛，涂抹后可以产生清爽和凉爽的感觉。

产品描述 : 一种清澈的浅棕色油，具有强烈的特征芳香气味。

配方

水杨酸甲酯 35% 薄荷醇 12.5% 樟脑 5% 桉树油 18%
薄荷油 7.5% 肉桂油 2% 丁香油 2% 白油 (up to) 100%

适应症 用于缓解肌肉和关节轻微疼痛的症状。

用法 将油涂抹在患处，每天揉搓 3 至 4 次。此药仅供外用。

禁忌

该产品可引起抽搐。它也禁止2岁以下儿童使用它。
年龄较大的儿童接受治疗时必须谨慎

警告和预防措施

将任何含有樟脑的产品放入儿童的鼻孔是很危险的。以这种方式应用的少量可能会导致立即崩溃。剧烈摩擦是不可取的。
本品含有水杨酸甲酯，涂抹或涂抹于皮肤时，可通过皮肤吸收进入血液。

与其他药物的相互作用

对于服用华法林的患者，因肌肉或关节疼痛而过度涂抹在皮肤上可能会增加出血的机会。

副作用 该产品可能会引起轻微的皮肤刺激或皮肤红斑，但是，这些是该产品的预期效果。由于搽剂是一种抗刺激剂和发红剂，因此刺激和红斑是预期的作用。

过量的症状和治疗

过量可能会导致皮肤刺激。

滥用

吞下药油可能会引起胃肠道症状，如呕吐和腹泻。治疗是对症的。大量误食后出现急性中毒，表现为恶心、呕吐、腹痛、头痛、眩晕、潮热、抽搐、呼吸抑制和昏迷。对有严重胃肠道或神经中毒症状的患者应进行观察和对症治疗。不要催吐。

怀孕期间使用

怀孕期间使用的安全性尚未确定。因此，应权衡产品的潜在益处与对母婴可能造成的风险。

储存条件

存放在 30°C 以下的干燥处。请将本品放在儿童接触不到的地方。

包装 5 毫升、10 毫升、30 毫升瓶

保质期 4 年

修订日期 29-7-2022

产品注册持有人/制造商:

SYARIKAT SAFETY MEDICAL

No. 48-2 & 50-2, Metro Perdana Timur 11
Jalan 13/34A, Kepong Entrepreneur's Park,
52100 吉隆坡 马来西亚

Torch Brand Liniment

Torch Brand Liniment is a common household medicine, useful and convenient to keep and carry with you. This is a special preparation which relieves pains and aches and can produce a refreshing and cooling feeling after application.

PRODUCT DESCRIPTION : A clear light brown oil with a strong characteristic aromatic odour.

FORMULA

Methyl Salicylate 35%	Menthol 12.5%	Camphor 5%	Eucalyptus Oil 18%
Peppermint Oil 7.5%	Cinnamon Oil 2%	Clove Oil 2%	White Oil Up To 100%

INDICATIONS For the symptomatic relief of minor aches and pains of muscle and joints.

ADMINISTRATION Apply the oil on affected part with rubbing 3 to 4 times daily. This medication is for external use only.

CONTRAINDICATIONS

This product can cause convulsion. It is also contraindicated in children under 2 years of age.
Caution must be exercised when older children are treated

WARNINGS AND PRECAUTIONS

It is dangerous to place any camphor containing product into the nostrils of children. A small amount applied this way may cause immediate collapse. Vigorous rubbing is not desirable. This product contains Methyl Salicylate and when applied or rub on the skin, can be absorbed through the skin into the blood.

INTERACTION WITH OTHER MEDICAMENTS

For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

SIDE EFFECTS This product can induce a mild irritation or erythema of the skin, however, these are the desired affects of the product. Since the liniment is a counterirritant and rubefacient, irritation and erythema are expected actions.

SYMPTOMS AND TREATMENT OF OVERDOSE

Over dosage may result in skin irritation.

Misuse

Swallowing of the medicated oil might cause gastrointestinal symptoms like vomiting and diarrhoea. Treatment is symptomatic. Acute poisoning was observed after significant accidental consumption, with nausea, vomiting, abdominal pain, and headache, vertigo, feeling hot flushing, convulsions, respiratory depression and coma. Patients with severe gastrointestinal or neurological symptoms of poisoning should be observed and treated symptomatically. Do not induce vomiting.

USE IN PREGNANCY

Safety for use in pregnancy has not established. Therefore, the potential benefits of the product should be weighed against the possible risks to the mother and child.

STORAGE CONDITION

Store in a dry place below 30°C. Keep out of reach of children.
Jauhkan daripada capaian kanak-kanak

PACK SIZE 5ml, 10ml, 30ml Bottle

SHELF LIFE 4 Years

DATE OF REVISION 29-7-2022

Product Registration Holder / Mfg by:
SYARIKAT SAFETY MEDICAL
No. 48-2 & 50-2, Metro Perdana Timur 11
Jalan 13/34A, Kepong Entrepreneur's Park
52100, Kuala Lumpur, Malaysia.