

UBAT MINYAK GELIGA - Liniment

Active Ingredient

Menthol Crystal 5.51% w/w, Camphor 9.2% w/w

Product Description

A light yellow clear liquid, odour characteristic.

Pharmacodynamics

Camphor : Is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, it becomes ideal for neuralgia and other painful area.

Menthol: Dilate the blood vessels causing a sensation of coldness, following by ab analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Pharmacokinetics

Camphor

Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.

Menthol

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Indication

For temporary relief of minor aches, pain of muscles and joints.

Recommended Dosage

Rub on affected area when necessary, 2 to 3 times a day.

Mode of Administration

Topical

Contraindications

- Hypersensitivity to any of the ingredients in the product.
- Product can cause convulsions. Contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated. Avoid direct application into nostrils

Warnings and Precautions

This product contains Camphor, it is contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated.

PRECAUTION:

It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

For external use only. Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Interactions with Other Medicaments

Has not been established, therefore the potential benefit of the product should be weighed against the possible risks to the patient.

Statement on Usage During Pregnancy and Lactation

Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse Effects / Undesirable Effects

This product can cause convulsions.

May cause hypersensitivity/allergic reactions in some individuals with sensitive skin.

Overdose and Treatment

Symptoms of camphor/menthol overdose

Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness, and coma; instant collapse in infants after the local application of menthol/camphor to their nostrils.

Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self-limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema, and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure, and death.

Treatment

The stomach should be emptied by gastric lavage or administration of oral activated charcoal

Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.

Any convulsions must be controlled first through supportive care including anticonvulsant therapy

Storage Conditions

Store below 30°C.

Dosage Forms and Packaging Available

Glass bottle , 5ml, 28ml, 60ml and 100ml

Name and Address of Manufacturer

Syarikat Wen Ken Drug Sdn Bhd

No 24, Lalan Lambak, Taman Johor, 81200 Johor Bahru Johor

Name and Address of Product Registration Holder

Syarikat MYP Sdn Bhd, 9-9A Jalan Petaling, Larkin Industrial Area , 80350, Johor Bahru, Johor

Date of Revision of Package Insert - AUG 2023