

Doxium® 500mg

Regulator of the capillary functions

Product Description

Hard gelatin capsule (cap: opaque dark green; body: opaque yellow) with a white powder fill.

Composition

Active principle: Calcium dobesilate monohydrate 500 mg.

Excipients: Maize starch, magnesium stearate. Capsule shell's composition: gelatin, yellow ferric oxide (E172), titanium dioxide (E171), Indigotin (E132).

Properties/Effects

Calcium dobesilate acts on the capillary walls by regulating its impaired physiological functions – increased permeability and decreased resistance. It increases erythrocyte flexibility, inhibits platelet hyperaggregation and, in diabetic retinopathy, it reduces plasma and blood hyperviscosity, thus improving blood rheological properties and tissue irrigation. These effects allow to correct capillary dysfunctions either of functional origin or caused by constitutional or acquired metabolic disorders.

Calcium dobesilate contributes to reduce oedema.

Pharmacokinetics

After oral administration of 500 mg of calcium dobesilate, its blood level is above 6 µg/ml between the 3rd and 10th hour, with a maximum (C_{max}) of 8 µg/ml on the average after 6 hours (t_{max}). Twenty four hours after intake blood level is about 3 µg/ml. The rate of protein-binding is 20 - 25%. In animals, calcium dobesilate does not cross the haematoencephalic or the placental barrier, but it is not known whether this is also the case in humans. Calcium dobesilate enters the maternal milk in very low quantities (0.4 µg/ml after intake of 1500 mg as observed in one study). Calcium dobesilate does not enter the enterohepatic cycle and is excreted mainly unchanged with only 10% being excreted as metabolites. About 50% of the orally

administered dose are eliminated in the first 24-hour urine and about 50% in the faeces. Plasma half-life is around 5 hours.

Kinetics in particular clinical situations

Kidney failure

The safety and efficacy of calcium dobesilate have not been studied in patients with kidney disorders. As the medication is excreted in the urine, care must be taken in the case of kidney failure. Consequently, the dose may be reduced for the administration of Doxium to such patients, in particular in patients with severe kidney failure requiring dialysis.

Liver failure

The safety and efficacy of calcium dobesilate have not been studied in patients with liver failure. Precautions should therefore be taken with the administration of Doxium to such patients. It is recommended that the benefit of the treatment be reassessed in the event of a significant increase in liver values.

Children

No trials have been conducted to study the use of calcium dobesilate in children.

Indications and usage

Microangiopathies, in particular diabetic retinopathy.

Clinical signs of chronic venous insufficiency in the lower limbs (pain, cramps, paresthesia, oedema, stasis dermatosis), as adjuvant in superficial thrombophlebitis.

Haemorrhoidal syndrome, micro-circulation disorders of arterio-venous origin.

Dosage

For adults only

Generally 500 to 1000 mg - 1 capsule once or twice a day - to be taken with the main meals.

Treatment duration, which is generally between a few weeks and several months, depends on the disease and its evolution.

Dosage should be adapted individually according to the severity of the case.

Doxium must be taken during or immediately after main meals, in order to minimise any gastric discomfort.

Route of Administration

Oral.

Contraindications

Hypersensitivity towards calcium dobesilate or to one of the excipients according to the composition.

Precautions

Dosage should be reduced in case of severe renal insufficiency requiring dialysis. In very rare cases (0.32/million patients), incidence estimated on the basis of spontaneous reports, the intake of calcium dobesilate may induce agranulocytosis, probably linked to a hyper-sensitivity reaction.

This condition may be expressed by symptoms such as high fever, oral cavity infections (tonsillitis), sore throat, anogenital inflammation and accompanying symptoms, that are often signs of an infection.

The patient should be told that by any sign of infection he/she must immediately inform his/her physician. In that case, it is essential to control without delay the blood formula and leucogram and to discontinue the treatment.

Doxium can trigger severe hypersensitivity reactions (anaphylactic shock or reaction). In the event of a hypersensitivity reaction, treatment must be stopped.

Pregnancy/Breast-feeding

Pregnancy category C:

Pregnancy

Data on the use of calcium dobesilate in pregnant women is limited. Studies conducted in and animals have not shown any direct or indirect harmful effect with regard to reproductive toxicity.

As a precaution, the medication should not be administered during pregnancy unless absolutely necessary.

Breast-feeding

Calcium dobesilate passes into breast milk in very low quantities

(0.4 µg/ml after taking 3 x 500 mg). The administration of Doxium is not indicated during breast-feeding. As a precaution, either treatment or breast-feeding should be suspended.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

No relevant studies have been conducted. Taking Doxium can cause undesirable effects such as nausea, headaches and fatigue. This is why care is recommended when driving and operating machines (see “Adverse effects

Adverse effects

Like all medicines, Doxium@500 can cause side effects, although not everybody gets them.
The adverse effects reported are presented by MedDRA system organ class and by frequency, as follows:listed below:

- Very common (≥1/10)
 - Common (≥1/100 to <1/10)
 - Uncommon (≥1/1,000 to <1/100)
 - Rare (≥1/10,000 to <1/1,000)
 - Very rare (<1/10,000)
 - Unknown frequency (cannot be estimated from the available data)
- Blood and lymphatic system disorders
Very rare: agranulocytosis (see section “warnings and precautions”)
Unknown: neutropenia, leukocytopenia
- Immune system disorders
Uncommon: hypersensitivity (rash, allergic dermatitis, pruritus, urticaria, facial oedema; see section “precautions”)
Very rare: anaphylactic reaction (see section “precautions”)

- Nervous system disorders
Common: headache
- Gastrointestinal disorders
Common: abdominal pain, nausea, diarrhoea, vomiting.
- Hepatobiliary disorders
Common: increased alanine aminotransferase

- Musculoskeletal and connective tissue disorders
Common: joint pain, muscle pain
- General disorders and administration site conditions
Uncommon: pyrexia, shivering, asthenia, fatigue
- These reactions are generally reversible when stopping treatment course.
In case of gastrointestinal disorders, the dosage should be reduced or the treatment temporarily withdrawn.
In case of skin reactions, fever, articular pain or change in blood formula, the treatment must be stopped and the treating physician informed as this may constitute hypersensitivity reactions.
If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

- Interactions**
At therapeutic doses, calcium dobesilate can interfere with creatinine enzyme measurements resulting in lower values than expected.
- During treatment with Doxium, the taking of a sample (e.g. a blood sample) required for laboratory tests must be done before the first administration of the medication in order to minimise any potential interaction between Doxium and the laboratory tests.

- Overdosage**
No cases of overdose have been reported and the clinical signs of overdose are not known. Overdose must be treated in accordance with standard medical practice.

- Particular remarks**
Incompatibilities
No incompatibility is known up to now.

- Information**
At therapeutic doses, calcium dobesilate may interfere with the assay of creatinine (too low values).

Special precautions for storage
Store below 30°C. Protect from heat and humidity.
The medication should not be used after the expiration date printed on the package together with the mention EXP.

Presentation
Box of 30 or 60 capsules.

This is a medicinal product

- The medicinal product is a product which affects your health, and its consumption contrary to instruction is dangerous for you.
- Follow strictly the doctor’s prescription, the method of use and instruction of the pharmacist who sold the medicinal product.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicaments out of reach of children

Manufacturer:
Corden Pharma Lisbon S.A.
Rua Da Industria, N 2,
Amadora,
2610-088 Portugal

Product Registration Holder:
Pharmaforte (M) Sdn Bhd
2, Jalan PJU 3/49
Sunway Damansara
47810 Petaling Jaya, Selangor, Malaysia

Updating of the information: February 2024

MY-DX51223

	Product: Doxium 500mg capsule Country: Malaysia
Item code: 2055405836	Version code: MY-DX5 1223
Dimensions: 297 x 190 mm	Date: 26-02-2024 v2
Cutter guide:	Laetus code: XXX
Colors:  Black	
Fonts:	