

**ZADIMIN SYRUP**

Banana flavoured plain syrup.

Content

Each 5ml contains Ketotifen hydrogen fumarate equivalent to ketotifen 1.0mg

Indications

Long term prevention of all forms of bronchial asthma, allergic bronchitis, asthmatic symptoms associated with hay fever. Prevention or treatment of multi system allergies, allergic rhinitis and allergic skin reactions.

Pharmacology

Ketotifen has anti allergic properties and has been used similarly to sodium cromoglycate in the prophylactic treatment of asthma. It also has the properties of an antihistamine. Ketotifen is absorbed from the gastrointestinal tract after oral administration. Together with its metabolites, it is excreted unchanged in the urine and feces.

Adverse effects

Increased appetite and weight gain have been reported. A reversible fall in the platelet count has been observed in a few patients receiving ketotifen concomitantly with oral antidiabetic agents and it has been suggested that this combination should therefore be avoided. Sedation and rarely dry mouth or slight dizziness may occur at the beginning of treatment but usually disappear spontaneously with continued medication. Occasionally, symptoms of CNS stimulation have been observed.

Precaution

Zadimin may potentiate the effects of CNS depressants, antihistamines and alcohol. Antiasthmatic drugs already in use should never be withdrawn abruptly when long term treatment with Zadimin is begun. This applies especially to systemic corticosteroids, because of the possible existence of adrenocortical insufficiency in steroid dependant patients: in such case, recovery of a normal pituitary adrenal response to stress may take up to one year. During the first few days of treatment with Zadimin the patients reactions may be impaired. Care should therefore be exercised when driving a vehicle, operating machinery, etc.

A reversible fall in the thrombocyte count in patients receiving Zadimin concomitantly when oral antidiabetic agents has been observed in rare cases. Thrombocyte counts should therefore be carried out in patients taking antidiabetics concomitantly.

Although there is no evidence of any teratogenic effect Zadimin should be given to pregnant and nursing women only under compelling circumstances.

Contraindications Nil.**Symptoms and treatment for overdosage and antidote**

Symptoms include drowsiness, confusion, dyspnoea, bradycardia or tachycardia, disorientation and convulsions. Gastric lavage is necessary and recovery within 12 hours after supportive treatment.

Dosage and Administration

For oral use

Adult dose

1-2 teaspoonfuls twice daily. In susceptible patients, start with ½ teaspoonfuls twice daily.

Children

More than 3 years : 1 (5ml) teaspoonfuls twice daily.

6 months - 3 years : ½ (2.5ml) teaspoonfuls twice daily.

Storage : Keep container tightly closed. Store in a cool, dry place below 30°C.
Protect from light.

Presentation : 120ml/ bottle

Shelf-life : 2 years

Manufactured & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by: 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.