

SWECON® TABLET

Swecon (Hydrotalcite) is a single antacid preparation which rapidly and consistently adjusts the acidity of the stomach to pH of 3-5. At this therapeutically optimal pH-range, excess gastric acid is quickly and lastingly bound to stop acid-induced pain.

Ingredient(s):

Each chewable tablet contains:
Magnesium Aluminium Hydroxide Carbonate Hydrate (Hydrotalcite)
..... 500mg

Pharmacodynamics & Pharmacokinetics:

Pharmacodynamics:

- a) Swecon acts very rapidly and consistently to adjust the acidity of the stomach to a pH of 3 ~ 5. At this therapeutically optimal pH-range excess gastric acid is bound and lastingly, and acid-induced pain is stopped.
- b) It lastingly inhibits pepsin activity and other components of gastric juice do not affect the antacid action.
- c) It acts locally and has no systemic effects. It is not absorbed and does not interfere with the acid-base balance. It has no laxative and no constipating effect and have been well tolerated.

Pharmacokinetics:

Swecon acid-binding rate is 100 % and acts very rapidly and at the same time persistently in the therapeutically optimal pH range between 3 and 5.

Indication(s):

Peptic ulcer (Gastro-duodenal ulcer), acute and chronic gastritis, hyperacidity.

Dosage and Administration:

To be chewed before swallowing
Usual adult dose is 2 tablets three-four times a day taken between meals or before sleep, depending on the age and severity of the case.

Route of Administration: To be taken orally.

Contraindication:

Contraindicated in patients who is hypersensitivity to any of its components.

Side Effect(s) / Adverse Reaction(s):

Occasionally: Diarrhea, soft stool, dry mouth and anorexia.

Precaution(s) / Warning(s):

Swecon should be given with caution to patients with renal disorder, cardiac disorder, diarrhea and hypermagnesemia. Absorption may be decreased, when oral tetracyclines are used concurrently with antacids. Patients should be advised not to take antacid within 1 to 2 hours after administering tetracycline.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description:

A round concave white tablet, one side is impressed with "SWECON" and the other side is impressed a circle with a dot at the center "



Shelf-Life:

4 years from the date of manufacture.

Packing(s):

Plastic bottle of 1000's (for export only).
Blister packing of 10's x 10, 10's x 120 and 10's x 100

The information contained in this leaflet is limited. Further information can be obtained from other sources including your doctor or pharmacist.

Item Code	: 204012 - 05 draft 18 Mar 2022
PM Code	: Ins.P SWT
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: L 139 - L 141 mm
CURSOR (ATTACHMENT 2)	: L 016 - L 021 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC

L
260mm
±
2mm

SWECON® TABLET

Bahan aktif:
Setiap tablet kunyah mengandungi:
Magnesium Aluminium Hydroxide Carbonate Hydrate (Hydrotalcite)
..... 500mg

Indikasi:
Ulser peptik, gastritis akut dan kronik, hiperasiditi.

Dos:
Kunyah sebelum telan.
2 tablet 3 hingga 4 kali sehari selepas makan dan sebelum tidur,
bergantung pada umur dan keamatan kes.

適胃康錠

成份:
每嚼錠中含:
Magnesium Aluminium Hydroxide Carbonate Hydrate (Hydrotalcite)
..... 500mg

適應症:
消化性潰瘍、急性胃炎、胃酸過多。

用法、用量:
咀嚼服用
成人 每次 2 粒 每日 3 ~ 4 次，於兩餐之間或睡前服用，
劑量和服用次數隨年齡和症狀調整。

Item Code	: 204012 - 05 draft 18 Mar 2022
PM Code	: Ins.P SWT
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: L 139 - L 141 mm
CURSOR (ATTACHMENT 2)	: L 016 - L 021 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC

L
260mm
±
2mm



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 18 Mar 2022

W
110mm ± 2mm