



PRIME MIXTURE POTASSIUM CITRATE

Description: Plain, lemon flavoured syrup.

Content:

Potassium citrate 1.5gm/5ml

Indications:

It is used principally for alkalinisation of urine.

Pharmacology:

Potassium citrate has the action of alkali citrate and is used principally to make the urine alkaline. It increases the alkali reserve and renders the urine less acid. It is used to make the urine alkaline in the treatment of infections of urinary tract and during treatment with some sulphonamides to prevent crystalluria.

Adverse reactions:

Confusion, irregular heartbeat, numbness or tingling in hands, feet, lips, shortness of breath, unexplained anxiety, unusual tiredness or weakness or heaviness of legs, bad taste, mood or mental changes and muscular pain or twitching.

Precaution:

It should be used with caution in pregnancy, breast-feeding, pediatrics and geriatrics.

Drug interactions

- Quinidine : Concurrent use with citrates may inhibit urinary excretion and prolong the duration of action of these medications.
- Antacids : Concurrent use with potassium citrates may result in systemic alkalosis.
- Antimuscarinics : Concurrent use with potassium citrates may increase the risk of gastro-intestinal irritation because of slowed gastro-intestinal transit time.
- Milk or other dairy products except butter. Concurrent use with citrates may increase the risk of calcium stones since alkaline urine may precipitate calcium.
- Digitalis glycosides : Concurrent use with potassium citrate may increase the risk of hyperkalemia in digitalized patients.
- Diuretics, potassium sparing.
- Low salt milk.
- Potassium containing medications.
- Salt substitutes : Concurrent use with potassium citrate may increase serum potassium concentrations, which may cause severe hyperkalemia and lead to cardiac arrest. Low salt milk may contain up to 60mEq of potassium/L and most salt substitute contain substantial amount of potassium.
- Laxatives : Concurrent administration with citrates may have an additive effect since potassium citrate may act as a saline laxative.
- Levodopa : Concurrent use with citrates may increase absorption of levodopa, especially in patient with slow gastric emptying.
- Methenamine : Concurrent use with citrates is not recommended because alkalinizing the urine may inhibit the effects of methenamine.
- Salicylates : Concurrent use with citrates may increase the urinary excretion and decrease the therapeutic effects of salicylates due to alkalization of the urine.

Contraindications:

It should not be used in patients with heart failure, myocardial damage, renal impairment or insufficiency.

Dosage and Administration

Usual adult dose : 10ml three times a day. Children : 5ml three times a day. To be taken well diluted with water.

Overdosage

For hyperkalemia

- Ascertaining food and medication containing potassium and any potassium sparing diuretics.
- Administering 10% dextrose injection containing 10 - 20 units of insulin/liter at a rate of 300 - 500ml of solution/hr.
- Correcting acidosis with intravenous sodium bicarbonate.
- Using exchange resins, hemodialysis or peritoneal dialysis.
- Observing caution when treating hyperkalemia in digitalized patients, since rapid reduction of serum potassium concentrations may reduce digitalis toxicity.

For metabolic alkalosis

- Controlling symptoms by patient rebreathing expired air into a paper bag or mask.
- Administering calcium gluconate injection if alkalosis is severe, to control tetany.

Storage conditions : Keep container tightly closed.

: Store in a cool, dry place, below 30°C. Protect from light.

Shelflife : 3 years

Presentation : 120ml/ bottle

Manufactured & : PRIME PHARMACEUTICAL SDN. BHD.

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah, 14000 Bukit Mertajam, Penang, Malaysia.

12/05/2017