

COPASTIN® FILM COATED TABLET 10mg

Copastin®, brand of Cloperastine Hydrochloride, an non-narcotic antitussive agent, is a cough suppressant which is used in the treatment of unproductive cough. Cloperastine Hydrochloride is an antitussive agent with an effect similar to that of codeine, and it also has some antihistaminic effect.

Ingredient(s):

Each tablet contains:

Cloperastine Hydrochloride 10mg

Pharmacodynamic:

1. Cloperastine Hydrochloride is known to reduce cough as a result of its central action, although the exact mechanism is still not entirely clear.
2. Cloperastine Hydrochloride has mild papaverine-like action and some antihistaminic effect.
3. Cloperastine hydrochloride is a centrally acting cough suppressant with some antihistamine action. It is a non opoid antitussive with an effect similar to that of codeine.
4. Cloperastine suppresses the cough reflex by a direct effect on the cough centre in the medulla of the brain. It exhibits the combination of a strong central antitussive effect, papaverine-like action and a mild antihistamanic action.
5. Copastin® Film Coated Tablet has no addictive and hypnotic effect and does not depress respiration.

Pharmacokinetic:

1. Cloperastine Hydrochloride is readily absorbed from the gastrointestinal tract, its effects occurring between 20 and 30 minutes and lasting for about 3 - 4 hours.
2. The effectiveness of Cloperastine Hydrochloride in patients with pathological cough has been demonstrated in controlled studies, where it was found to be about the equal of codeine. Unlike codeine, it rarely produces drowsiness or gastrointestinal disturbances. Its toxicity is quite low.

Indication(s):

For unproductive cough induced by common cold, acute and chronic bronchitis.

Dosage and Administration:

For adults: 1 – 2 tablets, three times daily.

To be used as directed by the physician.

Route of Administration:

Oral

Contraindication(s):

Known hypersensitivity to this product.

Warning and Precautions:

Use with caution in condition characterized by the production of bronchial secretions, such as cystic fibrosis, as these may result in sputum retention and the development of pneumonia.

Interactions with Other Medicaments:

None has been mentioned.

Pregnancy and Lactation:

No adverse effect on the fetus has been reported. However, Cloperastine should be given to women who suspect pregnancy or to pregnant women and nursing mothers only when clearly needed.

Side Effect(s) / Adverse Reaction(s):

Anorexia, nausea, dry mouth and drowsiness may occur.

Symptoms and Treatment for Overdosage and Antidote(s):

Symptoms of overdosage includes severe drowsiness, depression, confusion, tiredness and nausea. There is no specific antidote for Cloperastine. Treatment of overdosage consists of emptying the stomach via induction of emesis or gastric lavage, administering activated charcoal, and monitoring and supporting vital function.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product Description:

A faint green colored film coated round tablet, one side impressed with a "C".

Dosage Forms and Packaging Available:

Plastic bottle of 500's and 1000's (for export only).

Blister packing of 10's x 100.

COPASTIN[®] TABLET BERSALUT FILEM

Bahan Aktif:

Setiap tablet mengandungi:

Cloperastine Hydrochloride 10mg

Indikasi:

Batuk kering yang disebabkan oleh selesema biasa, bronkitis akut dan kronik dan lain-lain.

Dos:

Dewasa: 1 – 2 tablet, 3 kali sehari.

Digunakan seperti yang diarahkan oleh doktor.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679