

DESCRIPTION:

GEL

Colour : Natural

Flavour : Apricot

CONTAINS:

Triamcinolone (as Acetonide) 0.1% w/w

PHARMACODYNAMICS:

Triamcinolone provides a synthetic corticosteroid with marked anti-inflammatory action. The synthetic analogues are primarily used for their potent anti-inflammatory effects in disorders of many organ systems. By using Triamcinolone-containing gel, the possibility of systemic effects arising from ingested medication is thereby reduced. Furthermore, Triamcinolone-containing gel provides a soothing and protective covering, which augments the effects of the steroid.

PHARMACOKINETICS:

The absorption and fate of Triamcinolone is similar to corticosteroids. When administered by topical application particularly under an occlusive dressing or when the skin is broken, or as a pulmonary aerosol inhalation or a rectal enema, sufficient corticosteroid may be absorbed to give systemic effects. Corticosteroids in the circulation are extensively bound to plasma proteins, mainly to globulin and less so to albumin. The corticosteroid-binding globulin has high affinity but low binding capacity, while the albumin has low affinity but large binding capacity. Only unbound corticosteroid has pharmacological effects or is metabolised. Corticosteroids are metabolised mainly in the liver but also in the kidney, and are excreted in the urine. Triamcinolone is reported to have a biological half-life in plasma of about 300 minutes. It is bound to plasma albumin to a much smaller extent than hydrocortisone.

INDICATIONS:

Indicated for adjunctive treatment and for temporary relief of symptoms associated with oral inflammatory lesions and ulcerative lesions resulting from trauma.

RECOMMENDED DOSAGE:

Apply a small dab of Triamcinolone-containing gel to the lesion. A larger quantity may be required for coverage of some lesions. For optimal results use only enough to coat the lesion with a thin film. The gel should be applied at bedtime to permit steroid contact with the lesion throughout the night. Depending on the severity of symptoms, it may be necessary to apply the preparation 2 or 3 times a day, preferably after meals.

CONTRAINDICATIONS:

The use of topical corticosteroids is contraindicated in hepatic lesions of known viral origin, e.g. herpes labialis, intra-oral lesions, e.g. primary herpetic gingival stomatitis and herpanginas.

Contraindicated in patients with history of hypersensitivity to any of its component and propylene glycol.

PRECAUTIONS/WARNINGS:

Patients with tuberculosis, peptic ulcer or diabetes mellitus should not be treated with corticosteroid preparations including Triamcinolone-containing gel, without the advice of the patient's physician.

Discontinue if local irritations develop.

Review treatment in 7 days if repair or regeneration of oral tissue has not occurred.

Immunisation procedures should not be undertaken in patients who are on corticosteroids, especially on high dose, because of possible hazards of neurological complications and a lack of antibody response.

PREGNANCY AND LACTATION:

The use of this drug in pregnancy, nursing mothers or women of childbearing potential requires that the possible benefits of the drug be weighed against the potential hazards to the mother and the embryo, foetus or nursing infant. Infants born of the mothers who have received substantial doses of corticosteroids during pregnancy should be carefully observed for signs of hypoadrenalism.

SIDE EFFECTS/ADVERSE REACTIONS:

Intolerance to Triamcinolone-containing gel is extremely rare. Among several hundred patients treated for mucosal lesions, only one significant reaction to the vehicle components occurred: A strongly allergic individual developed giant hives on several occasions after application of the gel. No topical reactions to the corticosteroid have been observed with Triamcinolone-containing gel applied in the mouth. Moreover, the small amounts release when the gel is used as recommended make systemic effects very unlikely. However, if any unusual symptoms, e.g. weakness or dizziness should occur, they should be called to the attention of the patient's physician.

SYMPTOM AND TREATMENT FOR OVERDOSAGE:

For external use, there are no symptoms of an acute overdose known.

PACKING/PACK SIZE(S):

Aluminium tube of 10 g.

JAUHI UBAT DARIPADA KANAK-KANAK

KEEP OUT OF REACH OF CHILDREN

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, MALAYSIA.

PI1PE D005-05

Latest Revision: Sep 2024