

DYNA CALCIUM TABLET

MAL19984056X

DESCRIPTION:

TABLET

Colour : White
Shape : Round, Biconvex and Scored
Coating : Film-coated

EACH TABLET CONTAINS:

Calcium Lactate 300 mg equivalent to Calcium 39 mg

INDICATION:

As health supplement.

RECOMMENDED DOSE:

Adult and Adolescent (more than 12 years): One tablet at 3 - 4 hours intervals as needed, up to 25 tablets a day as required.

CONTRAINDICATION:

Contraindicated in patients with:

- Hypercalcemia
- Hypercalciuria
- Renal calculi, or ventricular fibrillation.
- Arcoïdosis (may potentiate hypercalcemia)

WARNING AND PRECAUTIONS:

Calcium salts should not be taken for prolonged periods without medical advice to patient with impaired renal function, cardiac disease, or sarcoidosis.

Calcium enhances the effects of digitalis on the heart and may precipitate digitalis intoxication; parenteral Calcium therapy is contraindicated in patients receiving cardiac glycosides. Calcium salts reduce the absorption of tetracyclines.

DRUG INTERACTIONS:

Alcohol or Caffeine (usually more than 8 cups of coffee a day), or tobacco (concurrent use of excessive amounts of these substances has been reported to decrease Calcium absorption).

Calcitonin

Concurrent use with Calcium supplements may antagonize the effect of calcitonin in the treatment of hypercalcemia; however when calcitonin is prescribed for osteoporosis or Paget's disease of the bone, Calcium intake should be generous to prevent hypocalcemia which might generate secondary hyperparathyroidism.

Gallium Nitrate

Concurrent use with Calcium supplements may antagonize the effect of Gallium Nitrate.

Tetracyclines, oral

Concurrent use with Calcium supplements may decrease absorption of tetracyclines because of possible formation of non-absorbable complexes and increase in intragastric pH; patients should be advised not to take Calcium supplements within 1 to 3 hours of taking tetracyclines.

Digitalis

Calcium enhances the effects of digitalis on the heart and may precipitate digitalis intoxication; parenteral Calcium therapy is contraindicated in patients receiving cardiac glycosides. Calcium salts reduce the absorption of tetracyclines.

Phenytoin

Concurrent use with Phenytoin reduces bioavailability of both Phenytoin and Calcium. Patients are advised not to take Calcium salts within 1 to 3 hours of taking Phenytoin.

SIDE EFFECTS/ADVERSE REACTIONS:

Side/adverse effects may be more likely to occur if oral Calcium supplements are taken in much larger doses than recommended (greater than 2000 to 2500 mg a day), if they are taken for a longer period of time or if they are taken by patients with renal function impairment or milk-alkali syndrome.

Excessive administration of Calcium salts leads to hypercalcaemia.

Abdominal pain, chalky taste and constipation.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Clinical feature:

Sign of potential overdosage if oral Calcium supplements are taken in much larger doses than recommended (greater than 2000 to 2500 mg) especially Calcium renal calculi or hypercalcemia.

For treatment of overdose:

A serum Calcium concentration exceeding 2.6 mol per liter (10.5 mg per 100 ml) is considered a hypercalcemia condition.

Withholding additional administration of Calcium and any other medications that may cause hypercalcemia usually resolves mild hypercalcemia in asymptomatic patients, when patient renal function is adequate.

PACKING/PACK SIZE(S):

Blister pack of 3x10's, 10x10's and 100x10's.

JAUHI DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN

RECOMMENDED DAILY ALLOWANCES

MINERAL	Unit	Adults and Children 4 or More Years of Age	Pregnant & Lactating Women
Calcium	g	1.0	1.3

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

PI1NT D001-03

Latest Revision: May 2022