

OTSU-NS 0.9% SODIUM CHLORIDE INTRAVENOUS INFUSION B.P.

COMPOSITION:

Each 500 mL contains:

Sodium Chloride 4.50 g Water for Injection
ad 500 mL

DESCRIPTION:

Clear, colorless, sterile and pyrogen free solution.

PHARMACOLOGY:

Sodium Chloride are the principle extracellular ions, their average serum concentrations are 142 and 103 mEq/L per liter (327 and 365 mg per 100 mL) respectively. The extracellular concentrations of both ions are much lower. The reduced cellular levels of sodium may be maintained by a secretory mechanism depending for its energy on cellular metabolism which moved potassium into the cell.

Sodium Chloride 0.9% is infused when sodium and water have been depleted in isotonic proportions. It may be used to maintain effective extracellular fluid volume and a stable circulation during and after surgery in patients with normal cardio vascular and renal function and postponing the need for blood transfusions in emergencies.

INDICATION:

- In general to overcome Sodium and Chloride deficiency in the blood, for example in case of dehydration and others.
- As a solvent for parenteral medicaments.

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION:

As required

SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTE (S) :

Signs and symptoms of excessive therapy are hypernatremia, eg. weakness, dehydration, dizziness, peripheral edema and pulmonary congestive (rales, edema)

TREATMENT:

Use of Sodium containing infusion should be ceased and other sodium intake controlled. In severe hypernatremia, dialysis may be indicated.

WARNING AND PRECAUTION:

- Careful administration for patients with heart failure, renal malfunction, peripheral and pulmonary edema, severe sepsis, pre-post trauma condition, children and the elderly.
- Do not use if bottle is leaking, solution is cloudy or contains foreign matter.

SIDE EFFECT & ADVERSE REACTION:

Parenteral administration could cause edema, if given in large amounts or the infusion rate exceeding the capacity of the blood circulation. This could happen to patients with malfunction of heart / kidney.

DRUG INTERACTION: No known data

PACKAGE:

Plastic bottle of 100 mL and 500 mL

SHELF LIFE:

5 years from the date of manufacturing

STORAGE:

Below 30°C (normal storage condition)

“Control Medicine/Ubat Terkawal”

ON MEDICAL PRESCRIPTION
ONLY



Manufactured by:
PT. Otsuka Indonesia
Jl. Sumber Waras No. 25
Lawang, Malang 65216, Indonesia

Product Registration Holder

Luen Wah Medical Co. Sdn. Bhd.
No. 43G & 43-1 F, The Highway Centre
Jalan 51/205, 46050 Petaling Jaya
Selangor Darul Ehsan, Malaysia