

P.I.

100mm

## OTSU-HM HARTMANN INTRAVENOUS INFUSION BP

### Compound Sodium Lactate Intravenous Infusion B.P.

#### COMPOSITION:

Each 500 mL contains:

|                       |           |                                   |                     |
|-----------------------|-----------|-----------------------------------|---------------------|
| Sodium Lactate        | 1.600 g   | Sodium                            | 131 mM/L            |
| Potassium Chloride    | 0.200 g   | Potassium                         | 5 mM/L              |
| Calcium Chloride 2H O | 0.135 g   | Calcium                           | 2 mM/L              |
| Sodium Chloride       | 3.000 g   |                                   |                     |
| Water for injections  | ad 500 mL | Bicarbonate (as Lactate) Chloride | 29 mM/L<br>111 mM/L |

#### SPECIFICATION:

Clear, colorless, sterile and pyrogen free solution. pH 5.0 - 7.0

Osmolarity 278 mOsm/L

#### INDICATION:

As a fluid and electrolyte replenisher, temporary blood volume increase, systemic alkalizer for acidosis followed by dehydration.

#### DOSAGE:

Intravenous administration, the dosage according to the necessity and patient condition individually.

#### SIDE EFFECT/ADVERSE REACTIONS:

No known data available

#### CONTRAINDICATION:

Patients with known hypersensitivity to component of the drug.

#### WARNING AND PRECAUTION:

- Administration of Sodium Lactate Intravenous Infusion carries the risk of producing lactic acidosis, particularly in seriously ill patients with poor tissue perfusion, heart failure and severe liver disease.
- Not for use in the treatment of lactic acidosis.
- Do not use if the bottle is leaking, solution is cloudy or contains foreign matter.

#### DRUG INTERACTION:

No Known data available

#### SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTE (S):

Signs and symptoms of excessive therapy are peripheral edema and pulmonary congestion (rales/edema and lactic acidosis).

#### PACKAGE:

Plastic bottle of 500 mL

#### STORAGE:

Store below 30°C

Keep medicine out of reach of children  
Jauh dari Kanak-kanak

Do not use if bottle is leaking, solution is cloudy or contains foreign matter.

“Control Medicine/Ubat Terkawal”

ON MEDICAL PRESCRIPTION ONLY



Manufactured by:  
PT. Otsuka Indonesia  
Jl. Sumber Waras No. 25  
Lawang, Malang 65216, Indonesia

Product Registration Holder:  
Luen Wah Medical Co Sdn Bhd  
No 53-2, 49-G, 49-1, The Highway Centre,  
Jalan 51/205, 46050 Petaling Jaya,  
Selangor Darul Ehsan, Malaysia

Revision Date: 15 July 2020

CILBSMLXXXXX

220 mm