

ET CREAM

(Antifungal - Anti-inflammatory - Antipruritic)

Active Ingredients:

Econazole Nitrate 1%w/w
Triamcinolone Acetonide 0.1%w/w
Preservative - Chlorocresol 0.1%w/w

Presentations:

Tube of 10g and box of 10g x 50 tubes.

Product Descriptions:

White smooth cream.

Pharmacology:

ET Cream is a combination of broad spectrum antimycotic, Econazole Nitrate and topical corticosteroid, Triamcinolone Acetonide with anti-inflammatory, antipruritic and antiallergic effects.

Econazole Nitrate is an imidazole antifungal agent which acts against dermatophytes (Trichophyton, Epidermophyton, Microsporum, Pityrosporum); pathogenic yeast (Candida albicans and other candida species) and moulds. The mechanism of action is by damaging the cell membrane, thus increasing the permeability to substances otherwise cannot penetrate into the intact cells, and therefore killing them. It is also active against G+ve bacteria, eg. Staphylococci, Streptococci, Bacillus anthracis and Bacteroides fragilis.

Triamcinolone Acetonide is a high potency glucocorticoid. It has a higher lipid-water distribution coefficient than Triamcinolone which favours penetration into the skin and is thus more suitable for topical use. It is employed for the treatment of various sensitive inflammatory and allergic dermatoses.

Triamcinolone Acetonide may be absorbed through the skin when applied topically, particularly under an occlusion dressing or when the skin is broken. Corticosteroids in the circulation are extensively bound to plasma protein, mainly to globulin and less so to albumin. The corticosteroid binding globulin has higher affinity but less binding capacity; while the albumin has low affinity but large binding capacity. Only unbound corticosteroid has pharmacological effects or is metabolised.

Metabolism occurs mainly in the liver, but also in the kidney and are excreted in the urine. Urinary excretion of 17 -OH corticoids is used as an index of adrenal function. Affinity may account for their increased potency compared with the natural corticosteroids. The absorption is not significant when econazole nitrate is applied to the skin.

Indications:

For the treatment of inflammatory dermatomycoses caused by dermatophytes, yeasts and moulds.

Directions of Use:

Apply the cream to the affected skin areas twice a day and rub in gently. Cleanse the affected area with warm water before application.

Treatment should if possible be limited to a period of 3 to 4 weeks, in order to prevent skin damage caused by corticoids (atrophy, telangiectasis, striae). Treatment of inflammatory mycoses should be continued until inflammation disappears. If necessary, the therapy can be carried on using Winderm Cream (Econazole Nitrate 1%w/w), until complete cure is obtained.

Symptoms and Treatments of Overdose:

Toxic effects result from the continued use of large doses of corticosteroids and from its withdrawal. Treatment is symptomatic; no specific antidotes available. Where possible, the dosage should be reduced or the drug withdrawn.

Contraindications:

Contraindicated in cases of tuberculous, luetic or virus infections (herpes, vaccinia, varicella) of the skin or varicose veins. Do not use if hypersensitive to imidazole or corticosteroids.

Precautions:

For external use only. Avoid in contact with eyes, nose and mouth. Not to be taken.

Avoid prolonged application to the face.

Avoid occlusive dressing to large areas of the body because of the increased risk of systemic absorption.

If symptoms persist, please consult a doctor.

Use in Pregnancy and Lactation:

Safety in pregnancy has not been established. Econazole should not be used in pregnant women or in those likely to become pregnant unless the expected benefit outweighs the potential risk.

It is not known whether Econazole is excreted in breast milk or whether it has a harmful effect on the new born. However as systemic absorption is low in humans, any excretion into breast milk is likely to be low.

Undesirable Effects:

Local reactions including slight burning sensation and irritation may occur.

Storage:

Store below 30°C. Protect from light. Keep cap or container tightly closed after opening.

Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

195101009677(175794-A)

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,

Pulau Pinang, Malaysia.

MAL19985553AZ

Date of Revision: 25 June 2025

A.EET-1.08

Size : 75mm x 115mm
■ Dark Blue (Pantone 2767U)