

ISORADIN CREAM

Isoradin is a water miscible white cream containing the active ingredient 1% isoconazole nitrate and 0.1% diflucortolone valerate. This cream is effective in the topical treatment of superficial mycosis of the skin caused by dermatophytes (trichophyton, Epidermophyton and Microsporum species), yeast and yeast-like fungi (including Candida albicans).

Ingredient(s):

Each gram contains:

Isoconazole Nitrate 10mg
Diflucortolone 21-valerate 1mg

Summary of Pharmacology (Pharmacodynamics and Pharmacokinetics):

Pharmacodynamics:

1. Isoconazole nitrate probably acts on the fungal cell wall membrane including permeability changes which alter the ionic macromolecular composition of the affected cells.
2. In vitro the spectrum of action of isoconazole nitrate includes the common dermatophytes, yeast and yeast-like fungi.
3. Diflucortolone valerate possesses excellent anti-inflammatory, antipruritic and anti-allergic properties. It is indicated to provide symptomatic relief of inflammation and/or pruritus associated with acute and chronic adrenocorticoid-responsive disorders.

Pharmacokinetics:

1. Isoconazole nitrate is not activated in the skin. Studies on percutaneous absorption indicate that over a 7 hours period, approximately less than 7% of the applied dose of isoconazole nitrate cream is absorbed from intact healthy skin.
2. Increased percutaneous absorption of diflucortolone valerate may occur when the skin or mucosa is abraded or inflamed, when body temperature is elevated, with prolonged use, or with extensive use.

Indication(s):

For the treatment of fungal infected eczema, inflammatory fungal infection, and infections due to fungi and gram positive bacteria.

Dosage and Administration:

Apply lightly to cover the affected area, twice daily. To avoid recurrence, treatment should be continued until clinical healing is achieved. To be used as directed by a physician.

Route of Administration:

Topical administration.

Contraindication(s):

1. Contraindicated in patients with known hypersensitivity to any of its components.
2. Should not be used for treatment of tuberculous lesions and viral diseases of the skin, nor should it be applied to varicose ulcers or to the eyes.

Precaution(s) / Warning(s):

Discontinue treatment if severe irritation or other signs of allergic reaction occur during use.

Pregnancy and Lactation:

Since safety of topical corticosteroid use in pregnant women has not been established, drug of this class should be used during pregnancy only if the potential benefit justifies the potential risk to fetus. Application to large areas of the body (about 10% and more) and/or for long periods of time (more than 4 weeks) during pregnancy must be avoided.

It is not known whether topical application of corticosteroid can result in sufficient systemic absorption to produce detectable quantities in breast milk. Topical steroid should not be applied to the breast area prior to breast feeding. When topical corticosteroid treatment is considered necessary for other parts of the body, both the amount applied and the length of treatment should be minimised.

Interactions with Other Medicament(s):

None has been reported.

Side Effect(s) / Adverse Reaction(s):

Isoradin is generally well tolerated. Infrequent adverse reactions include smarting or burning sensations, irritation, and maceration.

The following reactions may occur when applied to large areas of the body (about 10% and more) and/or for long periods of time (more than 4 weeks):

local concomitant symptoms such as atrophy of the skin, telangiectasia, striae, acneform changes of the skin, perioral dermatitis, increased growth of body hair (hypertrichosis), and systemic effects of the corticoid due to absorption.

Symptoms And Treatment For Overdosage, And Antidote(s):

There is no specific antidote for its overdose; therefore, management of the patients consists of symptomatic and supportive therapy.

Effects on Ability to Drive and Use Machine:

No effects on ability to drive and use machines have been observed in patients treated with topical isoconazole nitrate and difluocortolone valerate.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in an air tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description:

A white to off-white cream.

Dosage Form and Packaging Available:

Aluminium tube of 10g, 20g, 10g x 50, 20g x 72 and 20g x 40.

ISORADIN KRIM

Bahan Aktif:

Setiap gram mengandungi:

Isoconazole Nitrate	10mg
Difluocortolone 21-valerate	1mg

Indikasi:

Untuk rawatan jangkitan ekzema fungi, jangkitan fungi inflamasi dan jangkitan yang disebabkan oleh fungi dan bakteria Gram-positif.

Dos:

Sapukan sedikit krim di bahagian yang terlibat dua kali sehari. Untuk mengelakkan perulangan, rawatan mesti diteruskan sampai sembuh.

淨黴 乳膏

成份：

每公克含：

Isoconazole Nitrate	10mg
Difluocortolone 21-valerate	1mg

適應症：

治療因黴菌感染所致的皮膚炎症。

如：發炎性癬症和感染性濕疹。

用法、用量：

早晚一次，塗抹於患部，必須治療至症狀消失為止以防復發。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679