

Neuromethyn

Capsule 500 μ g

(Mecobalamin)

Neuromethyn is a mecobalamin preparation as a treatment for peripheral neuropathies. Neuromethyn contains mecobalamin, a vitamin B₁₂-coenzyme that occurs in the blood and the cerebrospinal fluid; it is taken up by nerve tissues more actively and extensively than other homologous of vitamin B₁₂. Biochemically, Neuromethyn accelerates the metabolic pathway of nucleic acids, proteins and lipids through its involvement in the transmethylation reaction; thus, it exerts a repairing effect on injured nerve tissues.

Clinically, Neuromethyn is useful for the treatment of numbness, pain and paralysis due to peripheral neuropathies such as diabetic neuropathy and polyneuritis.

Pharmacology (Summary of Pharmacodynamic & Pharmacokinetics): Vitamin B₁₂ occurs in the body mainly as mecobalamin and as adenosylcobalamin and hydroxocobalamin. Mecobalamin acts as coenzymes in nucleic acid synthesis. Vitamin B₁₂ binds to intrinsic factor, and is actively absorbed from the gastrointestinal tract by passive diffusion. Vitamin B₁₂ is extensively bound to specific plasma proteins(transcobalamins). It is stored in the liver, excreted in the bile, and undergoes extensive enterohepatic recycling. Vitamin B₁₂ diffuses across the placenta.

■ Description

Neuromethyn capsules are red in colour. They come in blisters of 10 capsules; 10 such blisters in a box.

■ Composition

Each capsule contains
Mecobalamin ----- 500 μ g

■ Indications

Peripheral neuropathies, megaloblastic anemia with Vit. B₁₂ deficiencies

■ Route of Administration

Oral administration

■ Recommended Dosage

Usually for adults, administer orally 3 capsules(1,500 mg of mecobalamin)in 3 divided doses per day. The dose may be adjusted according to the age of patients and severity of symptoms.

■ Symptoms & Treatment of overdosage

None has been reported so far as excessive intakes of most water-soluble vitamins have little effect due to their rapid excretion in urine.

■ Contraindications

No known contraindications have been reported so far.

■ Drug Interactions

No known drug interactions have been reported so far.

■ Warnings and Precautions

General Precautions

Neuromethyn should not be administered for prolonged periods(months) to patients who show no clinical response.

Mecobalamin contains cobalt and may cause sensitivity reactions in individuals with cobalt allergy.

■ Adverse Effect / Undesirable Effect

Digestive

Symptoms such as anorexia, nausea or diarrhea may infrequently occur.

Dermatologic

Skin rash may rarely occur.

■ Others

Prolonged use of larger doses of Neuromethyn not recommended for patients whose occupation requires handling mercury or its compounds.

■ Usage during Pregnancy and Lactation

Safety of Mecobalamin during pregnancy and lactating has not been established.

Thus, it should be used with caution in pregnancy and lactating.

■ Storage

Store in a cool dark place with tight, light-resistant container below 30°C.
Keep medicines out of reach of children

■ Shelf life

3 years from manufacturing date.

■ Date of Revision

Aug. 2024

MAL19988462X

SIN10917P



Manufactured by
SAMCHUNDANG PHARM.CO.,LTD.
Seoul, Korea

PRODUCT REGISTRATION HOLDER

The Zyfas Medical Co.

No. 15, Jalan Mega 1/5, Taman Perindustrian Nusa
Cemerlang, 79200 Iskandar Puteri, Johor.

Zyfas Pharma Pte. Ltd.

102F Pasir Panjang Road #06-10
Citilink Warehouse Complex Singapore 118530