

REXOM™

WINACOL SYRUP FOR CHILDREN

Name and Strength of Active Ingredient(s):

Guaifenesin	50mg/5ml
Preservative:	
Sodium Benzoate	0.1% w/v

Product Description

Pink syrup solution with strawberry flavour.

Pharmacodynamics

Guaifenesin reduces the viscosity of tenacious sputum and is used as an expectorant.

Pharmacokinetics

Guaifenesin is readily absorbed from the gastrointestinal tract. It is metabolised and excreted in the urine.

Indications

An expectorant for the relief of chesty coughs.

Recommended Dose

Children:

Age: 2 - 6 years old: Take 1 - 2 teaspoons (5 - 10ml) every 4 hours as required.

Age: 6 - 12 years old: Take 2 - 4 teaspoons (10 - 20ml) every 4 hours as required.

Do not use in children less than 2 years of age. Please consult your doctor before use.

Please use with caution and consult doctor's/pharmacist's advice in children 2 to 6 years of age.

Route of Administration

Oral

Contraindications

Hypersensitivity to the active substance.

Warning and Precautions

Ask a doctor before use if you suffer from a chronic cough, if you have asthma or are suffering from an acute asthma attack.

Stop use and ask a healthcare professional if your cough lasts for more than 5 days, comes back, or is accompanied by a fever, rash, or persistent headache.

Do not take with a cough suppressant.

Do not give this medicine with any other cough or cold medicines.

Interactions with Other Medicaments

There are no clinically significant drug interactions.

Other forms of interaction:

If urine is collected within 24 hours of a dose of guaifenesin a metabolite of guaifenesin may cause a colour interference with laboratory determinations of urinary 5-hydroxy-indoleacetic acid (5-HIAA) and vanillylmandelic acid (VMA).

Pregnancy & Lactation

The safety of this product during pregnancy and lactation has not been established.

However guaifenesin is not considered to constitute a hazard during pregnancy since there is no evidence of an association with foetal malformations. There is no data available on use of guaifenesin during breast feeding.

Side Effects

The following side effects may be associated with the use of guaifenesin:

Gastrointestinal Disorders: Nausea, vomiting, gastrointestinal discomfort.

Immune System Disorders: Hypersensitivity reactions.

Symptoms and Treatment of Overdose

Overdosage may give rise to nausea and vomiting. Treatment need only be symptomatic and supportive.

Effects on ability to drive and use machine

No adverse effects known.

Storage condition

Store below 30°C. Protect from light. Keep container tightly closed.

Dosage Forms

Oral Solution

Pack Size

60ml bottle

Registration Number

MAL19990611X

Manufactured by & Product Registration Holder

KCK PHARMACEUTICAL INDUSTRIES SDN. BHD.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

Date of Revision :

10 June 2021

PI0019-2