

SPAMODYN TABLET 40MG

DESCRIPTION:

Light yellow coloured round tablet with score on one side

COMPOSITION:

Each tablet contains Drotaverine hydrochloride 40mg

ACTIONS & PHARMACOLOGY:

Drotaverine is a musculotropic antispasmodic drug acting directly on smooth muscle cells by adsorption into the cell surface. The action does not involve the autonomic nervous system. It produces smooth muscle relaxation particularly at the site of the spasm. Drotaverine inhibits the phosphodiesterase enzyme, which leads to an increase of the CAMP level producing smooth muscle cell relaxation. Drotaverine also inhibits the initial calcium uptake of the cell. Drotaverine can safely be administered to patients suffering from glaucoma and prostatic hypertrophy, when anticholinergic antispasmodics are contraindicated. Drotaverine is well absorbed from the G.I.T. Drotaverine binds less to serum proteins than papaverine or perparin.

INDICATIONS:

For the relief of spasm of smooth muscles in the gastrointestinal tract.

CONTRAINDICATIONS:

Contraindicated in patients sensitive to drotaverine.

Concurrent use with levodopa is contraindicated as the antiparkinson effect may be decreased.

SIDE EFFECTS/ADVERSE REACTION:

Generally well-tolerated. Side-effects may occur rarely, such as gastrointestinal disturbances, nausea, constipation, headache, dizziness, vertigo, insomnia, palpitation and orthostatic hypotension.

WARNING AND PRECAUTIONS:

Exercise caution in patients with renal/hepatic/cardiac dysfunction.

PREGNANCY AND LACTATION:

Studies in animals have not revealed any teratogenic effect. However, as with most drugs, the use of drotaverine should be avoided during pregnancy and lactation unless essential.

DRUG INTERACTIONS:

The antiparkinson effect of levodopa may decrease when administered concomitantly with drotaverine.

DOSAGE & ADMINISTRATION:

Adults

By mouth 3 to 6 tablets per day in divided doses of 1 or 2 tablets

Children over six years

By mouth 2 to 5 tablets per day in divided doses of 1 tablet

Children between one to six years

By mouth 2 to 3 tablets per day in divided doses of ½ to 1 tablet

ROUTE OF ADMINISTRATION:

Oral administration

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE:

In therapeutic doses, drotaverine has no effect on ability to drive or operate machinery during oral treatment. Patients should be instructed that if they experience vertigo, they should avoid potentially hazardous tasks such as driving or operating machines

OVERDOSAGE & TREATMENT:

In case of overdosage, extended effects of the adverse reactions occur, including reduced gastric motility, hypotension and drowsiness. Treatment is symptomatic and supportive.

PRESENTATION:

Blister of 10 tablets. Box of 1, 3, 10, 50 and 100 blisters.

STORAGE CONDITIONS:

Store in a cool dry place below 30°C.

Protect from light.

Keep medicines out of reach of children.

MANUFACTURED BY:

Noripharma Sdn Bhd 200701034604 (792633-A)

Lot 5030, Jalan teratai.

5½ Mile Off Jalan Meru.

41050 Klang

Selangor Darul Ehsan., Malaysia.

DATE OF REVISION:

August 2021