

SVC Insert
110(L) x 260(H)mm
Draft (1) 03 May 2019

HEPAVITE® CAPSULE

Ingredient(s):

Each capsule contains :

Silymarin	70 mg
Thiamine Mononitrate (Vitamin B ₁)	3.9 mg
(e.q to Thiamine HCl	4 mg)
Riboflavin (Vitamin B ₂)	4 mg
Pyridoxine HCl (Vitamin B ₆)	4 mg
Nicotinamide (Vitamin B ₃)	12 mg
Calcium Pantothenate (Vitamin B ₅)	8 mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Silymarin is extracted from the seeds of the milk thistle (*Silybum marianum* Gaertn), a species known as an efficacious medicinal plant. A number of animal experiment were carried out and this drug was shown to protect liver damage from various hepatotoxic drugs. The therapeutic action of Silymarin is in stabilizing the liver cell membrane as well as the biomembranes of the organelles against toxic influences.

The vitamins of B-Complex are components of enzymes and participate as actual biocatalysts in many reaction cycles which occur in the liver cells. They accelerate the reparation of damaged liver parenchyma, and thus promote the organ's detoxifying action. Moreover, liver cells damaged through exogenous or endogenous toxins have a reduced accumulation capacity for vitamin B due to cell membranes no longer fully functioning. Therefore administration of the B-Complex vitamins can compensate such deficiencies.

Silymarin undergoes enterohepatic circulation and is mainly excreted via bile. Its major component, silybin, is excreted mainly by the kidney, although metabolites, particularly sulfate and glucuronide conjugates, are also present in bile. The total amount of silybin excreted in bile is about 20% - 40% of the ingested dose. Only 3% - 7% of the total dose is excreted renally.

Indication(s):

As an adjuvant in the treatment of the following diseases such as chronic liver disease, liver cirrhosis, fatty liver and intoxication by hepatotoxic substance.

Dosage and Administration:

1-2 capsules depending on the severity of the case, three times daily after meal.

Route of administration:

Oral administration

Contraindication(s):

Contraindicated in patients hypersensitive to any of its components.

Side Effect(s) / Adverse Reaction(s):

Silymarin and water-soluble vitamins seldom cause toxicity in persons with normal renal function.

Precaution(s) and Warning(s):

Problems have not been documented with intake of normal daily requirements. Studies have not been done in pregnant women, therefore Hepavite should be given to a pregnant woman only if clearly needed.

Drug Interaction(s):

Interactions with other drugs are not known.

Symptoms And Treatment For Overdosage and Antidote(s):

There is no specific antidote for its overdose. In case of overdosage, discontinue medication, treat symptomatically, and initiate supportive measures.

Shelf-life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 25°C. Protect from light and moisture.

Product Description and Packing(s):

A size #2 deep red cap and body hard gelatin capsule, containing yellow powder.

Blister Packing of 10's x 2, 10's x 10 and 10's x 100.

KAPSUL HEPAVITE®

Bahan Aktif:

Setiap kapsul mengandungi:

Silymarin	70 mg
Thiamine Mononitrate (Vitamin B ₁)	3.9 mg
(e.q to Thiamine HCl)	4 mg
Riboflavin (Vitamin B ₂)	4 mg
Pyridoxine HCl (Vitamin B ₆)	4 mg
Nicotinamide (Vitamin B ₃)	12 mg
Calcium Pantothenate (Vitamin B ₅)	8 mg

Indikasi:

Sebagai adjuvan untuk rawatan penyakit - penyakit seperti penyakit hati kronik, sirosis hati, hepar lemak, dan intoksikasi oleh bahan hepatotoksik.

Dos:

Bergantung kepada ketenatan kes, 1-2 kapsul tiga kali sehari, diambil selepas makan.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)
 Lot 3, 5 & 7, Jalan P/7, Section 13,
 Kawasan Perindustrian Bandar Baru Bangi,
 43000 Kajang, Selangor Darul Ehsan, Malaysia.
 Ordering Line: 1 800 88 3027
 Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 03 May 2019