
REXOM WINACOL SYRUP FOR ADULT

Name and Strength of Active Ingredients:

Guaifenesin.....100mg/5ml

Product Description

Dark Brown syrup solution with blackcurrant flavour.

Pharmacodynamics

Guaifenesin reduces the viscosity of tenacious sputum and is used as an expectorant.

Pharmacokinetics

Guaifenesin is readily absorbed from the gastrointestinal tract. It is metabolised and excreted in the urine.

Indications

An expectorant for the relief of chesty coughs.

Recommended Dose

Adults: 12 years and above:

Take 2-4 teaspoons (10-20ml) every 4 hours as required.

Route of Administration

Oral

Contraindications

Hypersensitivity to the active substance

Warning and Precautions

Not recommended for children under 12 years. Do not exceed the stated dose. Ask a doctor before use if you suffer from a chronic cough, if you have asthma or are suffering from an acute asthma attack and stop use and ask a healthcare professional if your cough lasts for more than 5 days, comes back, or is accompanied by a fever, rash or persistent headache. Do not take with a cough suppressant.

Interactions with Other Medicaments

There are no clinically significant interactions. If urine is collected within 24 hours of a dose of guaifenesin a metabolite of guaifenesin may cause a colour interference with laboratory determinations of urinary 5-hydroxyindoleacetic acid (5-HIAA) and vanillylmandelic acid (VMA).

Pregnancy & Lactation

The safety of this product during pregnancy and lactation has not been established. However guaiphenesin is not considered to constitute a hazard during pregnancy since there is no evidence of an association with foetal malformations. There is no data available on use of guaiphenesin during breast feeding.

Side Effects

Gastrointestinal disorders: nausea, vomiting/gastrointestinal discomfort.

Immune system disorder: hypersensitivity reactions, including anaphylaxis.

Symptoms and Treatment of Overdose

Overdosage may give rise to nausea and vomiting. Treatment need only be symptomatic and supportive.

Storage condition

Keep container tightly closed. Protect from heat and light. Store below 30°C.

Dosage Forms and Packaging Available: 60ml HDPE Bottle

Date of Revision: 22/5/2021

Registration Number: MAL19991909X

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.