

ELOSTEN-T CREAM

DESCRIPTION:

A white to off white, smooth homogenous mass.

COMPOSITION:

Each gm of the cream contains:

Econazole Nitrate (Micronised) 10 mg

Triamcinolone Acetonide (Micronised) 1 mg

ACTIONS & PHARMACOLOGY:

ELOSTEN-T is a combination of econazole nitrate and triamcinolone acetonide. Econazole nitrate is a broad spectrum antimycotic that acts against dermatophytes, yeasts and moulds. It is also active against gram-positive bacteria. Triamcinolone acetonide develops marked anti-inflammatory, antipruritic and antiallergic effects. Systemic absorption following topical application to normal skin is extremely low.

INDICATIONS:

Eczema unresponsive to less potent corticosteroids, inflammatory dermatoses with mycotic and or bacterial infection. Inflammatory dermatomycoses caused by dermatophytes, yeasts and moulds. Mycotic infections in the body folds.

CONTRAINDICATIONS:

Contraindicated in patients who are sensitive to topical econazole and triamcinolone. Contraindicated in tuberculous, luetic or virus infections (herpes, vaccinia, varicellae) of the skin.

SIDE EFFECTS:

ELOSTEN-T is well tolerated. Irritation, burning, itching, redness or other signs or irritation not present before therapy are some of the isolated reactions reported. Hypersensitivity was observed in very rare cases only. Discontinuation of therapy usually results in cessation of the symptoms.

WARNING & PRECAUTIONS:

Caution should be exercised when this preparation is administered to pregnant and nursing women. Avoid prolonged application to thin skin and to the face. Use of topical antifungals may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication. To reduce the possibility of recurrence, comply with full course of therapy. Avoid contact with the eyes. Long-term continuous steroid therapy should be avoided since adrenal suppression can occur, particularly when infants or children are treated or when occlusive dressings are applied. The preparation should be avoided in treating infants under 1 year or if necessary used with great care for short periods.

PREGNANCY AND LACTATION:

Topical administration of corticosteroids to pregnant animals can cause foetal abnormalities. The relevance of this finding to human beings has not been established. Nevertheless, the product should not be administered during pregnancy unless the possible risks involved have been carefully considered.

DRUG INTERACTIONS:

Interactions Econazole is a known inhibitor CYP3A4/2C9. However, due to the limited systemic availability after cutaneous application, clinically relevant interactions are unlikely to occur, but have been reported for oral anticoagulants. In patients taking oral anticoagulants, such as warfarin or acenocoumarol, caution should be exercised and the anticoagulant effect should be monitored.

DOSAGE & ADMINISTRATION:

Apply the cream to the affected skin areas twice a day (morning and evening) and rub in gently. Continue therapy until complete cure is obtained.

ROUTE OF ADMINISTRATION:

Topical Administration

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

None known.

OVERDOSAGE & TREATMENT:

Acute overdosage with topical application of the drug is unlikely. If irritation or sensitivity reactions occur, use of the medication should be discontinued. Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects eg. reversible HPA axis suppression, Cushing's syndrome, hyperglycaemia and glucosuria.

Treatment: Symptomatic, with the drug slowly withdrawn.

PRESENTATION:

Aluminium tubes of 5g, 10g, 15g, and 50g.

STORAGE CONDITION:

Store in a dry place below 30°C.

Keep the containers tightly closed.

Protect from light and freezing.

For external use only.

Keep medicine out of reach of children.

MANUFACTURED BY:

Noripharma Sdn. Bhd 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Miles Off Jalan Meru,

41050 Klang, Selangor Darul Ehsan, Malaysia.

DATE OF REVISION:

April 2021