

GELIGA AROMA -

Active Ingredient

Menthol Crystal 10% w/w,

Product Description

Light yellow clear liquid, odour characteristic.

Pharmacodynamics

Menthol : Dilate the blood vessels causing a sensation of coldness, following by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Pharmacokinetics

Menthol

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Indication

For temporary relief of minor aches, pains of muscle and joints

Recommended Dosage

Adults & children above 2 years old :
Apply few drops to the affected area
then rub or massage gently, 2 to 3 times a day,
when necessary

Mode of Administration

Topical

Contraindications

- Hypersensitivity to any of the ingredients in the product.
- Contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated. Avoid direct application into nostrils.

Warnings and Precautions

Keep all medicine away from children
Jauhkan daripada capaian kanak-kanak.

Interactions with Other Medicaments

Has not been established, therefore the potential benefit of the product should

be weighed against the possible risks to the patient.

Statement on Usage During Pregnancy and Lactation

Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child

Adverse Effects / Undesirable Effects

May cause hypersensitivity/allergic reactions in some individuals with sensitive skin

Overdose and Treatment

Symptoms of menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness, and coma; instant collapse in infants after the local application of menthol to their nostrils.

Treatment

The stomach should be emptied by gastric lavage or administration of oral activated charcoal
Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.
Any convulsions must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions

Store below 30°C

Dosage Forms and Packaging Available

External liquid, 5ml and 28ml in glass bottle

Name and Address of Manufacturer

Syarikat Wen Ken Drug Sdn Bhd
No 24, Jalan Lambak, Taman Johor, 81200,
Johor Bahru, Johor

Name and Address of Product Registration Holder

Syarikat MYP Sdn Bhd
9-9A Jalan Petaling, Larkin Industrial Estate, 80350
Johor Bahru, Johor

Date of Revision of Package Insert

- September 2025