



REMATRIL CAPSULES

Description: Red op./Cream op. capsules

Content

Each capsule contains:

Glucosamine sulphate sodium chloride 314.0mg
(eq. to glucosamine sulphate 250mg)

Indications:

As adjuvant therapy for osteoarthritis.

Pharmacology:

Glucosamine is an aminomonosaccharide, particularly effective in all forms of degenerative osteoarticular diseases e.g. arthrosis and osteoarthritis, both subacute and chronic, with disappearance or reduction of articular pains, improvement of articular functions and inhibition or regression of the degenerative process. Glucosamine possesses arthrotrophic metabolic activity by stimulating the anabolic metabolism of osteocartilaginous tissues, through the stimulation of the biosynthesis of the mucopolysaccharides (which are essential components of the cartilage substances) and of the bone mesenchymal tissues, and improvement of the bone calcium uptake.

Lubricant activity through improvement of synovial fluid viscosity by increasing synovial fluid production. After oral administration, only 26% is absorbed, with onset of action about 2-3 weeks for osteoarthritis. Bioavailability is low due to its first pass hepatic metabolism. Following oral administration, glucosamine is extensively metabolised in the liver. It is rapidly desulfated and metabolised into smaller molecules, and ultimately to carbon dioxide, water and urea. Glucosamine is not protein-bound, but rather incorporates into the plasma proteins (primarily globulin). Rapid diffusion into most tissues also occurs, including articular tissues. Urinary excretion accounts for 10% of an oral dose, about 11% is eliminated via the faeces, presumably as an unabsorbed drug. The elimination half-life of glucosamine which is incorporated into the plasma proteins is about 68 hours after oral doses. Part of a dose of glucosamine sulfate is eliminated as carbon dioxide via expired air in the lungs.

Adverse effects:

Cardiovascular: Peripheral edema, tachycardia were reported in a few patients following a larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central nervous system: Drowsiness, headache, insomnia has been observed rarely during therapy (less than 1%)

Gastrointestinal: Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation heartburn and anorexia have been described rarely during oral therapy with glucosamine.

SKIN:

Skin reaction such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Precautions:

Glucosamine sulphate is a causal therapy and the therapeutic effect is only seen after 1 week. Therefore, it is advisable to include an antiinflammatory drug (in case of severe pains) during the first days of treatment with glucosamine.

Derived From Seafood

Contraindications:

It is contraindicated in patients hypersensitive to glucosamine.

Dosage and Administration:

Light to moderate symptoms - 2 capsules twice daily for at least 6 weeks before meals.

Severe symptoms - 2 capsules three times daily before meals.

Capsules should be taken 15 mins. before each meal. Treatment should be repeated every other 6 months or less according to medical prescription.

Storage : Keep container tightly closed. Store in a cool, dry place below 30°C.
Protect from light.

Packing : 100's

Shelflife : 3 years

Product registration holder : Prime Pharmaceutical Sdn. Bhd.

& Manufacturer

1505, Lrg Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.