

**USER INFORMATION
READ CAREFULLY**

**INFUSOL® HS
SODIUM CHLORIDE 0.45% w/v INTRAVENOUS INFUSION BP**

Composition:

Sodium Chloride BP 4.5g/L

Electrolytes:	mmol/L	mEq/L
Sodium Ion (Na ⁺)	77	77
Chloride Ion (Cl ⁻)	77	77
Osmolarity :	154 mOsm/L	

Product Description:

Sodium Chloride 0.45% Intravenous Infusion BP is a hypotonic solution containing 77 mmol each of Sodium and Chloride per litre and supplies physiologically free water. The solution is a clear and colourless solution

Pharmacodynamic:

It is the principal salt involved in maintaining osmotic tension of the blood and tissues. Such change influences the movement of fluids and diffusion of salt in cellular tissues.

Normal osmolarity of the extracellular fluid range between 280 to 300mOsm/L; it is primarily a function of sodium and its accompanying ions, chloride and bicarbonate.

Sodium chloride is the principal salt involved in maintenance of plasma tonicity. One grams of sodium chloride provides 17.1 mEq of sodium and 17.1 mEq of chloride.

Pharmacokinetics:

Sodium Chloride is readily absorbed in the body. It is present in all types of body fluids but primarily in extracellular fluid. The osmotic equilibrium is maintained by excretion in urine.

Bioavailability: As the solution is directly administered into the blood, it is 100% available

Indication:

- Hypertonic dehydration.
- Light sodium and chloride depletion.
- Hypochloremic alkalosis.
- Vehicle solution for supplementary medication.

Recommended Dosage:

Unless otherwise prescribed: 1000mL/day

Drop Rate: 120-180 drops/min corresponding to 360-540mL/hr.

Route of administration: Intravenous (IV)

Contraindications:

- Hyperhydration
- Oedema
- Hypotonic dehydration
- Acidosis situations

Warnings and Precautions:

Excessive administration of sodium chloride causes hypernatraemia, the most serious effect of which is dehydration of internal organs, especially the brain. Excess chloride in the body may cause a loss of bicarbonate with an acidifying effect.

Interactions with Other Medicaments:

Compatibility of the solution should be checked before used.

Statement on usage during pregnancy and lactation:

As advised by the physician.

Adverse Effects/Undesirable Effects:

General adverse effects of excess sodium in the body include nausea, vomiting, diarrhoea, abdominal cramps, thirst, reduced salivation and lachrymation, sweating, fever, tachycardia, hypertension, renal failure, peripheral and pulmonary oedema, respiratory arrest, headache, dizziness, restlessness, irritability, weakness, muscular twitching and rigidity, convulsions, coma and death.

Reactions due to solution or technique of administration: Febrile response; local tenderness; abscess; tissue necroses or infection at injection site; venous thrombosis or phlebitis extending from injection site; extravasation; hypervolaemia.

If an adverse reaction occurs, discontinue the infusion, evaluate the patient, institute appropriate countermeasures and save the remainder of the fluid for examination.

Overdose and Treatment:

Hypernatraemia requires the use of sodium – free fluids and the cessation of excessive sodium intake. Very occasionally dialysis has been needed in severe hypernatraemia.

Iso-osmotic overload is managed by sodium and water restriction plus measures to increase renal sodium and water loss such as ‘loop diuretics’ (eg. Furosemide) or, in specific circumstances, antimineralocorticoid agents.

Cautions:

The compatibility of any additives to this solution should be checked before use. Single dose container. Discard all unused contents. Do not use if leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used.

Shelf Life:

5 years from manufacturing date in proposed storage condition in LDPE Bottle.

3 years from manufacturing date in proposed storage condition in PVC Medical Collapsible Bag.

Do not use after expiry.

Storage condition:

Do not store above 30°C.

Dosage form and packaging available:

100mL x 20 LDPE plastic bottle per carton

250mL x 20 LDPE plastic bottle per carton

500mL x 20 LDPE plastic bottle per carton

500mL x 10 LDPE plastic bottle per carton

500mL x 20 PVC Medical plastic collapsible bag per carton

Manufacturer / Product Registration Holder:



AIN MEDICARE SDN.BHD.



Lot 4933, 4934 & PT2464, Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA.

HANDLING INSTRUCTION

AIN MEDICARE INTRAVENOUS INFUSION FLUID IN LOW DENSITY POLYETHYLENE (LDPE) BOTTLE



The bottle is made from injectable grade polyethylene.

PARTS OF THE PLASTIC BOTTLE



Top view after pull ring cap is removed. Please refer whichever applicable design.


INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG



AIN MEDICARE INTRAVENOUS INFUSION FLUID IN PVC MEDICAL COLLAPSIBLE BAG

The Intravenous solution in medibag is packed and sterilized in an outerwrap.

PARTS OF THE COLLAPSIBLE BAG



INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG

