

**USER INFORMATION
READ CAREFULLY**

**INFUSOL® D5
DEXTROSE 5% w/v INTRAVENOUS INFUSION BP**

Composition:

The solution contains
Dextrose Anhydrous BP 50 g/L
Osmolarity 278 mOsm/L
Caloric Value : Approx. 837 kJ/L = 200 kcal/L

Product Description:

The solution is a colorless solution.

Pharmacodynamic:

Glucose provides energy.
Glucose when given intravenously with water metabolises into glycogen leaving water to be absorbed to correct fluid imbalance.
Glucose increases the movements of electrolytes across the semi-permeable membrane.

Pharmacokinetics:

Glucose is rapidly absorbed from the gastrointestinal tract. It is metabolised to carbon dioxide and water with the release of approx. 4 calories/gram of energy. It is readily metabolised. It may decelerate body protein and nitrogen losses, promote glycogen deposition and decrease/prevent ketosis in sufficient dose. As the product is administered through intravenous route it is 100 % bioavailable.

Indication:

Energy supply
Hypertonic dehydration
Vehicle solution for supplementary medication
Hypoglycaemia

Recommended Dosage:

Adult patients

According to individual requirements:
Dextrose 5% : up to 40mL/kg of body weight per day.
Drop rate (for patients with approximately 70kg of body weight)
Dextrose 5% : up to 120 drops/min or 360mL/h.

Pediatric patients

Mean requirements/kg of body weight and per day.
1st year of life : 8 - 15g dextrose.
2nd year of life : 12 - 15g dextrose.
3rd - 5th year of life : 12g dextrose.
6th - 10th year of life : 10g dextrose.

Drop rate

Dextrose 5% : up to 180 drops/kg of body weight and per hour.

Route of administration: Intravenous (IV)

Glucose solutions with a concentration greater than 5% are hypertonic and should be administered by slow intravenous infusion via a central vein.

Contraindications:

This solution is contraindicated in patients presenting with:
Diabetes mellitus (with the exception of hypoglycaemic conditions).
Glucose intolerance, hypotonic dehydration if lacking electrolytes are not replaced.
Over hydration.
Hypokalaemia.

Warnings and Precautions:

Blood glucose, serum electrolytes and water balance should be monitored regularly. Electrolytes are to be supplemented as required. The compatibility of any additives to this solution should be checked before use. Dextrose solutions should not be administered through the same infusion set through which also blood has or may be given because of the risk of pseudoagglutination.
This is a single dose container. Discard all unused contents. Do not use if leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used.

Interactions with Other Medicaments:

Compatibility of additive should be checked before used.

Statement on usage during pregnancy and lactation:

As advised by the physician.

Adverse Effects/Undesirable Effects:

Dextrose Injection and particularly hypertonic Dextrose Injection may have low pH, and such solution when injected may irritate the venous intima near the site of the injection to cause thrombophlebitis.
Hyperglycaemia and renal losses may occur in case of reduced glucose tolerance. These manifestations are normally prevented by reducing the dosage and/or giving insulin.
Enhanced bilirubin and lactate levels may be found if the recommended dosage is exceeded.

Overdose and Treatment:

Dextrose solutions I.V. can cause fluid overload resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary oedema.
Hyperglycaemia and glucosuria may occur if rate of administration is too fast or in case metabolic insufficiency. To minimise these conditions, reduce the infusion rate, monitor blood and urine glucose; if necessary, stop administration of dextrose or/and administer insulin.

Storage condition: Do not store above 30°C.

Shelf Life:

5 years from manufacturing date in the proposed storage condition for LDPE plastic bottle.
Do not use after expiry.

Dosage form and packaging available:

100mL x 20 LDPE plastic bottle per carton
250mL x 20 LDPE plastic bottle per carton
500mL x 20 LDPE plastic bottle per carton
1000mL x 10 LDPE plastic bottle per carton



Manufacturer / Product Registration Holder:




AIN MEDICARE SDN.BHD.
Lot 4933, 4934 & PT2464, Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA.

HANDLING INSTRUCTION

AIN MEDICARE INTRAVENOUS INFUSION FLUID IN LOW DENSITY POLYETHYLENE (LDPE) BOTTLE
The bottle is made from injectable grade polyethylene.
PARTS OF THE PLASTIC BOTTLE



Top view after pull ring cap is removed. Please refer whichever applicable design.
INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG

 Design 1	1. Before use check for any solid particle, growth, turbidity or leakage. Discard the product if particle, growth, turbidity or leakage is found.	 Design 2
 Design 1	2. Design 1: Pull the cap's ring until completely removed. Design 2: Pull the aluminium seal until completely removed by centre line.	 Design 2
 Design 1	3. Additives may be injected through injection site (Refer to Design 1 or 2 whichever applicable) on the rubber. Mix the solution thoroughly.	 Design 2
 Design 1	4. Hold bottle down firmly and insert a spike of administration set into the spike site for Design 1 or for Design 2, pull the other aluminium seal until completely removed before inserting a spike of administration set. <i>(Spike site is meant for spiking of the administration set and is only for single use)</i>	 Design 2