

**USER INFORMATION
READ CAREFULLY**

**INFUSOL® RG
RINGER'S INJECTION**

Product Description:

INFUSOL® RG is a full electrolyte solution for replacement of extracellular fluid and it is a colourless solution.

Composition:

The solution contains

Sodium Chloride BP	8.60 g/L
Potassium Chloride BP	0.30 g/L
Calcium Chloride BP	0.33 g/L
Osmolarity:	309.08 mOsm/L

Electrolytes:

Sodium Ion (Na ⁺)	147.16 mmol/L
Potassium Ion (K ⁺)	4.02 mmol/L
Calcium Ion (Ca ⁺⁺)	2.24 mmol/L
Chloride Ion (Cl ⁻)	155.66 mmol/L

Pharmacodynamic/Pharmacokinetics:

Sodium is the principle cation in the extracellular fluid and is responsible for maintenance of the extracellular fluid volume and osmolarity. Sodium Chloride is well absorbed from the gastro intestinal tract. Sodium is predominantly excreted by the kidney, but there is extensive renal reabsorption.

Calcium is involved in the maintenance of normal muscle and nerve function, is essential for normal cardiac function, and is essential to the clotting of blood. Potassium is the principle cation of intracellular fluid and is intimately involved in cell function and metabolism. It is essential for carbohydrate metabolism and glycogen storage and for protein synthesis. It is intimately involved in transmembrane potential and has profound effects on muscle, including heart muscle.

Indication:

Treats dehydration from reduced water intake, vomiting or diarrhoea. Treats mild alkalosis. Treats hyponatremia.

Recommended Dosage:

Average Dose: 1000mL/day

Drop Rate: 120-180 drops/min corresponding to 360-540mL/hr.

Route of administration: Intravenous (IV)

Contraindications:

Addison's disease

Severe potassium or calcium deficiency

Warnings and Precautions:

Sodium salts should be used cautiously in patients with cardiac failure, hypertension, impaired renal function, peripheral and pulmonary oedema and in toxemia of pregnancy

Statement on usage during pregnancy and lactation: As advised by the physician

Adverse Effects/Undesirable Effects:

Hypernatraemia, where the symptoms may include restlessness, weakness, thirst, reduced salivation and lachrymation, swollen tongue, flushing of skin, pyrexia, dizziness, headache, oliguria, hypotension, tachycardia, delirium, hyperpnoea, and respiratory arrest. Another is retention of sodium and water - isotonic retention or isosmotic expansion because the plasma concentration is unchanged, which leads to the accumulation of extracellular fluid (oedema).

Overdose and Treatment:

Hypernatraemia requires the use of sodium-free fluids and the cessation of excessive sodium intake. Very occasionally dialysis has been needed in severe hypernatraemia.

Isosmotic overload is managed by sodium and water restriction plus measures to increase renal sodium and water loss such as loop diuretics (eg. furosemide) or, in specific circumstances, antimineralocorticoid agents.

Cautions:

Single dose container. Discard all unused contents. Do not use if leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used.

Discard any portion not used when the seal is first broken. Before use check for minute leaks by squeezing bag firmly. Discard if leakage is detected.

Incompatibilities

Shelf Life:

5 years from manufacturing date in the proposed storage condition for LDPE Bottle.

3 years from manufacturing date in the proposed storage condition for PVC Medical Collapsible Bag.

Do not use after expiry.

Storage Conditions:

Do not store above 30°C.

Dosage form and packaging available:

250mL x 20 LDPE plastic bottle per carton

500mL x 20 LDPE plastic bottle per carton

1000mL x 10 LDPE plastic bottle per carton

500mL x 20 PVC Medical plastic collapsible bag per carton

1000mL x 10 PVC Medical plastic collapsible bag per carton

Manufacturer / Product Registration Holder:



AIN MEDICARE SDN.BHD.



Lot 4933, 4934 & PT2464, Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA.

HANDLING INSTRUCTION

AIN MEDICARE INTRAVENOUS INFUSION FLUID IN LOW DENSITY POLYETHYLENE (LDPE) BOTTLE





The bottle is made from injectable grade polyethylene.

PARTS OF THE PLASTIC BOTTLE



Top view after pull ring cap is removed. Please refer whichever applicable design.


INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG



AIN MEDICARE INTRAVENOUS INFUSION FLUID IN PVC MEDICAL COLLAPSIBLE BAG

The Intravenous solution in medibag is packed and sterilized in an outerwrap.

PARTS OF THE COLLAPSIBLE BAG



INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG

