

Jl. Dreded No. 34, Bogor 16132, Telp. 0251-315714 (hunting), Fax. 0251-336883

Materi Desain	Leaflet Regro 5% 80ml (41281-04)
Bharcode	-
Ukuran	177 x 126 mm
Klien	Actavis
Warna yang digunakan	 Cyan  Magenta  Yellow  Black
Bahan yang digunakan	
Finishing	-
Desain / Tanggal	
Desain oleh	



NAME AND STRENGTH OF ACTIVE SUBSTANCE: Minoxidil 5% w/v

PRODUCT DESCRIPTION: REGRO is a clear colourless to pale yellow mixture

PHARMACODYNAMICS: Following topical application, **REGRO** is poorly absorbed from normal intact skin, with only about 1-2% of the dosage ultimately reaching the systemic circulation.

PHARMACOKINETICS: It contains minoxidil which when applied topically has been shown to stimulate hair growth in individuals with alopecia androgenetica (male pattern baldness) for both male and female. The effectiveness of minoxidil can only be expected after 4 or more months of application on the affected area.

INDICATION: REGRO is indicated for the treatment of male pattern baldness for both male and female. Observations of patients suggest that the rate of balding progression may be slowed by the use of this product. Onset and degree of hair growth stimulation vary among patients. An inverse relationship may exist between duration of follicular dormancy and favourable response to treatment.

RECOMMENDED DOSAGE AND ADMINISTRATION: Apply 7 sprays (1ml) of **REGRO** two times a day, preferably morning and night, to the areas of baldness or hair loss. This amount should be used regardless of the size of the bald area. Do not exceed two doses (14 sprays or 2ml) per day. Massage scalp gently with fingertips after each spray. Wash hands thoroughly after use.

CONTRAINDICATIONS: REGRO is contraindicated in patients with a history of hypersensitivity to minoxidil, alcohol and propylene glycol.

PREGNANCY AND LACTATION: REGRO should not be used during pregnancy or lactation.

SIDE EFFECTS/ADVERSE REACTIONS: Local adverse reactions include scaling, dry skin, erythema/flushing, dermatitis, burning sensation and rash. Other infrequent side effects include allergic reactions, allergic rhinitis, facial swelling, seborrhea, headache, light headedness, vertigo, weakness, dizziness, edema, eye irritation and visual disturbances. Rarely reported adverse reactions include alopecia, hair abnormalities, hypotension, syncope, palpitations, pulse rate change, chest pain, shortness of breath, hepatitis and kidney calculi.

PRECAUTIONS/WARNING: Application of **REGRO** will lead to some absorption of minoxidil from the skin which may cause some systemic effects such as tachycardia, angina, oedema or potentiation of orthostatic hypotension produced by guanethidine. In the event of systemic side effects, application of the drug should be discontinued. If **REGRO** accidentally comes into contact with the eye, mucous membrane or abraded skin, the area should be rinsed with plenty of cool water. Inhalation of the spray mist should be avoided. It should not be used in conjunction with other topical corticosteroids, retinoids and petrolatum or agents that are known to enhance cutaneous drug absorption. Safety and effectiveness of minoxidil in patients under 18 years and over 65 years have not been established. It is for external use only. Do not apply to areas of the body other than the scalp.

OVERDOSAGE: Accidental ingestion may produce systemic effects related to the vasodilatory action of **REGRO**. Signs and symptoms of drug overdosage would most likely be cardiovascular effects associated with fluid retention, lowered blood pressure or tachycardia. Fluid retention can be managed with appropriate diuretic therapy. Tachycardia can be controlled by the administration of a beta-adrenergic blocking agent.

DRUG INTERACTION: No known drug interactions. Though not clinically demonstrated, there exists a possibility of potentiating orthostatic hypotension in patients concurrently taking peripheral vasodilators.

INCOMPATIBILITY: Not reported.

AVAILABILITY: Minoxidil 5% w/v X 80ml (with sprayer)

STORAGE: Keep below 30°C.

DATE OF REVISION: November 2021

For further information, please consult your physician or pharmacist.

Manufactured by:
PT Actavis Indonesia
Jl. Raya Bogor Km. 28, Kelurahan Pekayon,
Kecamatan Pasar Rebo
Kota Jakarta Timur, Provinsi DKI Jakarta, Indonesia



W
Actavis

AAAN8793

41281-04