

ELOSTEN CREAM

DESCRIPTION:

White smooth homogenous mass.

COMPOSITION:

Each gm of the cream contains

Econazole Nitrate (Micronised) 10mg

Preservative: Chlorocresol 0.1%w/w

ACTIONS & PHARMACOLOGY:

Econazole nitrate is a broad spectrum antifungal agent. Fungistatic; may be fungicidal, depending on concentration. Inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell wall membrane and altering its permeability. As a result, loss of essential intracellular elements may occur. Also inhibits biosynthesis of triglycerides and phospholipids by fungi. In addition inhibits oxidative and peroxidative enzyme activity, resulting in intracellular buildup of toxic concentrations of hydrogen peroxide, which may contribute to deterioration of subcellular organelles and cellular necrosis. In *Candida albicans*, inhibits transformation of blastospores into invasive mycelial form. It also has some activity against gram-positive bacteria. After topical application to the skin of normal subjects, systemic absorption of econazole nitrate is extremely low. Although most of the applied drug remains on the skin surface, drug concentrations were found in the stratum corneum which, by far, exceeded the minimum inhibitory concentration for dermatophytes. Inhibitory concentrations were achieved in the epidermis and as deep as the middle region of the dermis. Less than 1% of the applied dose was recovered in the urine and faeces.

INDICATIONS:

Elostén cream is indicated as a primary agent in the topical treatment of Tinea corporis (ring worm of the body) Tinea cruris (ring worm of the groin) or Tinea pedis (ring worm of the foot; athlete's foot) caused by *Trichophyton rubrum*, *T. mentagrophytes*, *T. tonsurans*, *Microsporum canis*, *M. audouini*, *M. gypseum*, and *Epidermophyton floccosum*. Elostén cream is also indicated as a primary agent in the topical treatment of cutaneous candidiasis (moniliasis) caused by *Candida* (*Monilia*) species and in the topical treatment of *Tinea versicolor*. Also indicated in the treatment of skin infections caused by gram positive pathogens.

CONTRAINDICATIONS:

In patients who are sensitive to econazole.

SIDE EFFECTS:

Side effects are less frequent. Hypersensitivity reactions which include burning, itching, stinging, redness, or other signs of irritation not present before therapy.

PRECAUTIONS:

Use of topical antifungals may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical use of the medication. To reduce the possibility of recurrence, *Candida* infections, Tinea cruris and Tinea corporis should be treated for at least 2 weeks and Tinea pedis should be treated for at least 1 month. When this medication is used in the treatment of candidiasis, occlusive dressings should be avoided since they provide conditions that favour growth of yeast and release of its irritating endotoxin. Use hygienic

measures to help cure infection or to help prevent reinfection. For Tinea cruris - avoid underwear that is tight-fitting or made from synthetic materials. For Tinea pedis - carefully drying feet, especially between toes, after bathing, avoid socks made from wool or synthetic materials; wearing well-ventilated shoes or sandals. Avoid contact with eyes.

Use in pregnancy and lactation

Caution should be exercised when econazole nitrate is administered to pregnant and nursing woman.

DOSAGE & ADMINISTRATION:

Apply to cover the affected areas once daily in patients with Tinea pedis, Tinea cruris, Tinea corporis and Tinea versicolor. Twice daily (morning and evening) in patients with cutaneous candidiasis.

OVERDOSAGE & TREATMENT:

Acute overdosage with topical econazole is unlikely. If irritation or sensitivity reactions occur, use of the medication should be discontinued. If accidental ingestion of large quantities of the product occurs, an appropriate method of gastric emptying may be used if considered desirable.

PRESENTATION:

Aluminium tubes of 5g, 10g, 15g and 50g

STORAGE:

Store in a cool dry place below 30°C. Keep the containers tightly closed.
Protect from light and freezing. For external use only.
Keep medicine out of reach of children.

Shelf - life: 2 years

MANUFACTURED BY:

Noripharma Sdn. Bhd
Lot 5030, Jalan Teratai.
5 ½ Miles off Jalan Meru,
41050 Klang,
Selangor Darul Ehsan