

Directions for use
Read carefully!



57/12623823/0717

B. Braun 0.33 % Sodium Chloride and 5 % Glucose Intravenous Infusion B.P.

Composition

Each 100 ml contains:

Sodium Chloride	0.33 g
Glucose Monohydrate for Parenteral Use	5.50 g
Water for Injections to	100 ml
Electrolytes:	mmol/l
Na ⁺	56
Cl ⁻	56
Caloric value	850 kJ/l $\triangle$ 200 kcal/l
Osmolarity:	390 mOsm/l

Characteristics

B. Braun 0.33% Sodium Chloride and 5% Glucose Intravenous Infusion B.P. is a light hypertonic solution containing 56 mmol each of sodium and chloride per litre and supplies physiologically free water.

Indications

Dehydration
Sodium and chloride depletion
Hypochloremic alkalosis
Vehicle solution for supplementary medication
Caloric supply

Dosage

Average dose: 1000 ml/day
Drop rate: up to 120-180 drops/ min corresponding to 360-540 ml/h

Contraindications

Hyperhydration, oedema
Hypernatremia
Hypotonic dehydration
Acidotic situation

Route of administration I.V.

Mechanisms of action

Glucose is absorbed from the gastrointestinal tract, and oxidised as a source of energy, or stored in the liver as glycogen. It is the only energy substrate which is directly, instantly and universally utilized by the body. Glucose is vital for the myocardium, brain and nerves.

Sodium Chloride is the principal salt involved in maintaining the osmotic tension of the blood and tissue. Changes in sodium and chloride levels change this osmotic tension and hence influence the movement of fluids and diffusion of salts in cellular tissue.

Symptoms and treatment of overdosage

If hyperglycaemia or glucosuria occur, the dosage and infusion rate should be reduced, or insulin may be administered.

Side effects/Adverse reactions

Hyperglycaemia and glucosuria may occur if glucose tolerance is impaired. These can be avoided by reducing the dosage and infusion rate or by administering insulin.

Precautions

Use with caution in patients with hypertension and diabetes mellitus. Blood glucose, serum electrolytes and water balance should be monitored regularly.

The compatibility of any additives to this solution should be checked before use.

Shelf life

The product must not be used beyond the expiry date stated on the label.

Storage

Do not store above 30 °C.

Presentation

500 ml plastic container

Approval for Printing

B. BRAUN Melsungen AG

Approved for Printing ☐

Approved for Printing
when corrected ☐

New draft required ☐

Date _____ Signature _____

Name in capital letters

schwarz

Dokument = 148 x 210 mm (DIN A5)
1 Seite

MY_57
Ecoflac Plus
B. Braun 0.33% Sodium Chloride
and 5% Glucose Intravenous Infusion B.P.
57/12623823/0717
Standort Penang

Lätus



4384

Font size: 9 pt.

G 170364

Product registration
holder and
manufactured by:
**B. Braun Medical
Industries S/B**
11900 Bayan Lepas,
Penang, Malaysia.

B. BRAUN

