

ABOUT BLACKMORES® BIO E 500

INDICATION: FOR VITAMIN E DEFICIENCY

Pharmacodynamics and pharmacokinetics:

- Vitamin E's most critical function is on the cell membrane. It acts as an anti oxidant, which prevents the destructive process caused by compounds called free radicals.
- Vitamin E converts the free radical into a less reactive, non-harmful form without the formation of another free radical in the process. This effect prevents oxidative damage to polyunsaturated fatty acids and the main structural components of cell membranes.

Pharmacokinetics:

- Absorption:** Vitamin E is a fat-soluble vitamin whose absorption from the gastrointestinal tract is dependent on normal pancreatic function and the

presence of bile. The amount absorbed varies between about 20% and 80% and appears to decrease as the dose is increased.

- Distribution:** Vitamin E enters the blood via the chylomicrons in the lymph and is bound to beta-lipoproteins. As it is critical for cell membrane protection, it is widely distributed to all tissues, and stored in adipose tissue.
- Metabolism:** Along with the distribution described, some vitamin E is metabolised in the liver to glucuronides, tocopheronic acid and its γ-lactone.
- Excretion:** Vitamin E is mainly excreted via the bile and only small amounts are eliminated in the urine. Vitamin E appears in breast milk but is poorly transferred across the placenta.

HOW TO USE:
Adults - Take 1 capsule a day with a meal. Not recommended for children under 12 years.

CONTRAINDICATIONS, SIDE EFFECTS, PRECAUTIONS AND WARNINGS:
Contraindication: Contraindicated in individuals who may be sensitive to any of the ingredients in the product.
Precautions & Warnings: Not suitable for use by pregnant or breastfeeding women. High dose vitamin E should be used with caution by those with cardiovascular disease or cancer of the head and neck.
Side Effects: Vitamin E is usually well tolerated. Large doses may cause diarrhoea, abdominal pain and other gastrointestinal disturbances, and have also been

reported to cause blurred vision, dizziness, fatigue and weakness. Large doses have been reported to increase bleeding tendency in vitamin K deficient patients such as those taking oral anticoagulants. However, it has also been suggested that it may increase the risk of thrombosis in some patients, such as those taking oestrogens. The clinical significance of these effects is not known.

- Medicine Interactions:** Various drugs may interfere with the absorption of vitamin E including cholestyramine, colestipol and orlistat. High doses of vitamin E may increase the effects of oral anticoagulants.
- Pregnancy and Lactation:** Not suitable for use by pregnant or breastfeeding women. The product contains 500 IU d-alpha-tocopherol. The Upper Level of intake for vitamin E during pregnancy and lactation

GLUE
15mm

ADHESIVE

GLUE
15mm

in Australia and New Zealand is 300 mg daily, which is equivalent to 447 IU d-alpha tocopherol.

- Symptoms of overdose and treatment:** Although side effects have been reported with extended intakes of 1000 to 2100 mg/day of vitamin E these effects are not severe and subside rapidly upon reducing the dosage or discontinuing the administration of vitamin E. Excess alpha-tocopherol, as well as forms of vitamin E not preferentially used, are probably excreted unchanged in bile. Serious ingestions require hydration if vomiting or diarrhoea is present. Oxygen and monitoring of vital signs are essential if potentially life-threatening manifestations are present. Consult a neurosurgeon if evidence of CNS haemorrhage is present. Vitamin K replacement through the oral or subcutaneous

route should decrease the risk of bleeding in patients who are taking anticoagulants or who have vitamin K deficiency.

- Storage instructions:** Keep out of reach of children. Store below 30°C in a dry place away from direct sunlight. Do not use the product beyond the expiry date printed on the pack.

ACTIVE INGREDIENT PER CAPSULE:

Each capsule contains **Vitamin E** (natural source) 500 IU (equivalent to d-alpha-tocopherol 335 mg).

Product description: A clear, unprinted, oblong shaped, soft gelatin capsule.

Presentation: Glass amber bottle with white cap containing 30 soft gelatin capsules.

ABOUT BLACKMORES:

Malaysia: Product registration holder: Blackmores (Malaysia) Sdn. Bhd., Unit No. 7-05-01, Level 5, Tower 7, UOA Business Park, No. 1, Jalan Pengaturcara U1/51A, Section U1, 40150 Shah Alam, Selangor, MALAYSIA. MAL20000314XCR. For more information ☎ **03-5022 8263.**

Singapore: Imported by: Blackmores (Singapore) Pte. Ltd. For more information ☎ **62253933.**

Distributed by: Zuellig Pharma Pte. Ltd., 15 Changi North Way, #01-01, SINGAPORE 498770.

Contract manufactured for Blackmores by: Catalent Australia Pty. Ltd., 217-221 Governor Road, Braeside VIC 3195 AUSTRALIA. **Packed by:** Blackmores Ltd., 20 Jubilee Avenue, Warriewood NSW 2102 AUSTRALIA.

Date of amendment: April 2025

72051K

BASE

BLACKMORES®

Label Size: 160 x 40 mm
Text Height: 1.5mm
NB: TGA Require 1.5mm min. text height

CMF Job No: 4932
New label Item Code: 43499
New Spec No: 72051K 543

NB: PHARMA CODE are last 3 digits of New Spec No.

PRODUCT NAME: BLACKMORES BIO E 500 (30) MY DOUBLE LABEL

Draft No: 3
Active Proof No: 1
Date: 19.08.25
EAN No:
93808422

100 ml Amber Glass Bottle

Colours used:

CYAN

MAGENTA

BLACK

PANTONE 7711

BLA030A–200

5
Colours

LABEL SPECIFICATION:

This label is a Blackmores standard label (MCC).

PRINTER DETAILS:

PRINTER MCC

SENT TO PRINT