



5/12622528/0919

Directions for use
Read carefully!

B. Braun 0.9% Sodium Chloride Injection B.P.

Composition

100 ml of solution contain

Active ingredients:

Sodium Chloride 0.9 g

Excipients:

Water for injections

Theoretical osmolarity: 308 mOsm/l

Titration acidity: < 0.3 mmol/l

pH: 4.5 - 7.0

Electrolyte concentrations:

Sodium 154 mmol/l

Chloride 154 mmol/l

Pharmaceutical form

Solution for injection

A clear and colourless solution

Pharmaco-therapeutic group

Vehicle solution

Pharmacodynamics

Mechanism of action and therapeutic effect

Sodium is the primary cation of the extracellular space and together with various anions, regulates the size of this. Sodium and potassium are the major mediators of bioelectric processes within the body.

The sodium content and the liquid metabolism of the body are closely coupled to each other. Each deviation of the plasma sodium concentration from the physiological one simultaneously affects the fluid status of the body.

An increase in the sodium content of the body also means reduction of the body's free water content independent of the serum osmolality.

A 0.9 per cent sodium chloride solution has the same osmolality as plasma. Administration of this solution primarily leads to a replenishment of the interstitial space which is about 2/3 of the entire extracellular space. Only 1/3 of the administered volume remains in the intravascular space. Therefore the haemodynamic effect of the solution is of short duration only.

Pharmacokinetics

Distribution

The total sodium content of the body is ca. 80 mmol/kg of which ca. 97 % is extracellular and ca. 3 % intracellular. The daily turnover is ca. 100 - 180 mmol (corresponding to 1.5 - 2.5 mmol/kg body weight).

Metabolism and elimination

The kidneys are the major regulator of the sodium and water balances. In co-operation with the hormonal control mechanisms (renin-angiotensin aldosterone system, antidiuretic hormone) and the hypothetical natriuretic hormone they are primarily responsible for keeping the volume of the extracellular space constant and regulating its fluid composition.

Chloride is exchanged for hydrogen carbonate in the tubule system and is, thus, involved in the regulation of the acid base balance.

Indications

Solvent for the preparation or dilution of parenterally administered drugs.

Contraindications

None

Precautions for use and special warnings

B. Braun 0.9 % w/v Sodium Chloride Injection should only be administered with caution in cases of

- hypernatraemia
- hyperchloraemia

Interactions

When mixing with other medicaments, possible incompatibilities should be considered.

Dosage

The quantity to be chosen depends on the desired concentration of the medicament to be dissolved.

Approval for Printing B|BRAUN Melsungen AG

Approved for Printing

☐

Approved for Printing
when corrected

☐

New draft required

☐

LRA:

Date

Signature

Name in capital letters

LMS:

Date

Signature

Name in capital letters

schwarz

Dokument = 148 x 210 mm

DIN A5

2 Seiten

MY__5

GIF (MPc, MPw)

5/12622528/0919

Production site: Berlin

Lätus: 7723



Font size: 9 pt.

G 160864

B|BRAUN





5/12622528/0919

Method of administration

Intravenous or subcutaneous injection.
For the use of this solution as solvent/diluent for compatible electrolyte concentrates or medications, the instructions for use relating to the medicament to be added must be observed.

Overdose

Symptoms

Overdose of B. Braun 0.9% Sodium Chloride solution may result in hypernatraemia, hyperchloraemia, hyperhydration, hyperosmolality of the serum, and metabolic acidosis.

Emergency treatment, antidotes

Immediate stop of administration, administration of diuretics with continuous monitoring of serum electrolytes, correction of electrolyte and acid-base imbalances.

Undesirable effects

None to be expected if the product is used according to directions.

Usage during pregnancy

Specific risks regarding the use during pregnancy are not known.

Storage

Do not store above 30 °C.

Expiry date

The product must not be used beyond the expiry date stated on the label.

Presentation

In plastic ampoules "Mini-Plasco®" and "Mini-Plasco® Connect" of 10 ml, 20 ml, in box of 20's.

Instructions for use / handling


The solution is supplied in single-dose containers. Discard unused contents.

Use contents immediately after opening the container.


Only to be used if solution is clear and the container undamaged.

Date of last revision: 11.2018

Mini-Plasco® Handling



Mini-Plasco® Connect Handling



B | BRAUN

Product registration holder:
B. Braun Medical Industries S/B
11900 Bayan Lepas, Penang
Malaysia

Manufacturer:
B. Braun Melsungen AG
Carl-Braun-Straße 1,
34212 Melsungen, Germany.

Production site:
B. Braun Melsungen AG
Mistelweg 2-6, 12357 Berlin,
Germany

