



PIMPLE GEL

DESCRIPTION

T3 Pimple Gel is an off-white to brownish gel with characteristic odour. It contains Resorcinol 2.0% w/w with Methyl Paraben 0.1% w/w and Propyl Paraben 0.05% w/w as preservatives.

INDICATIONS

Treat acne and pimple.

CONTRAINDICATIONS

Resorcinol is a mild irritant and may result in skin sensitization.

ADVERSE EFFECT

Resorcinol is a mild irritant and may result in skin sensitization.

PRECAUTIONS/WARNING

For external use only. Avoid contact with eyes. If excessive discomfort occurs, discontinue use. Resorcinol may produce hyperpigmentation in those with dark skins and may darken light-coloured hair.

INTERACTION WITH OTHER MEDICATIONS

Unknown

RECOMMENDED DOSAGE

At the first sign of pimples, apply T3 Pimple Gel directly onto affected spots after cleansing thoroughly with facial cleanser. Apply 3 times a day.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTES

It should not be applied to large areas of the body for prolonged period, or in high concentrations especially in children as it is absorbed through intact as well as broken skin and may interfere with thyroid function or produce methaemoglobinaemia.

PACKING

15g aluminium tube.

STORAGE

Keep container well closed. Protect from strong light. Store below 30°C. For external use only. Keep out of reach of children. Recommended shelf-life : 3 years.

Product Holder & Manufacturer:

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohd. 6,
Bandar Sultan Suleiman,
42000 Port Klang, MALAYSIA.

Revision Date: November 2017



PIMPLE GEL

说明

T3 暗疮凝胶是一种米白色至棕色凝胶，具独特香味。这种凝胶含有 2% w/w Resorcinol 及 0.1% w/w Methyl Paraben 和 0.05% w/w Propyl Paraben 作为防腐剂。

指示

治疗粉刺和暗疮。

禁忌象征

Resorcinol 是温和刺激物，可能会造成皮肤过敏。

副作用

Resorcinol 是温和刺激物，可能会造成皮肤过敏。

注意/警告

只限外敷，避免和眼睛接触。若有非常不适的感觉，即刻停止使用。Resorcinol 可能会在深肤色上形成染色，或使浅色的发色变深。

与其它药物的相互作用

未知。

建议用量

在暗疮长出的初期开始使用。在使用洁面霜清洗脸部后，可直接在暗疮部位涂上 T3 暗疮凝胶。每天使用 3 次。

过量使用的症状与治疗及解毒方法

切勿涂抹全身，长期涂抹或涂得太浓，特别是小孩，因为完好的肌肤和受伤的肌肤都会吸收这种药物，并且可能干扰甲状腺功能，或造成变性血红素贫血。

包装

15 克铝管包装。

存放

将瓶盖关紧，避免强光直晒。存放于摄氏 30 度以下的室内。只限外敷。远离孩童伸手可及之处。建议存放年限：3 年。

产品持有人及制造商

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohd. 6,
Bandar Sultan Suleiman,
42000 Port Klang, MALAYSIA.

修订日期：2017 年 11 月