



● SWKD
○ WKMS
○ WKMM
○ HU
○ HU (KK)

File Name: Povidone Iodine 30ml_Leaflet 251120
Received : 09.10.2018
Amended : 25.11.2020
Output : 25.11.2020
Request : Shuying
Designer : Syafiq
Opened : 165w x185h MM
Closed : -
Finished: -
Colour : 1C
K



Three Legs Povidone Iodine Povidone Iodine Cap Kaki Tiga 三脚标优碘

Composition: Povidone Iodine 10%w/v

Product Description: A dark brown colour with a slight characteristic odour solution.

Indication: For the treatment of wounds, cuts, abrasions and minor skin infections.

Direction for use: Apply twice daily to the affected area.

Route of Administration: Topical

Contraindication: Hypersensitivity to Povidone Iodine

Warning and Precaution: Application of Povidone Iodine to large areas of broken skin should be avoided as excessive absorption of iodine may occur.

When redness or swelling occurs due to hypersensitivity, stop application and consult doctor.

For external use only.

Drug Interaction: Use with concurrent lithium therapy has been shown to exhibit additive hypothyroidic effects. Absorption of iodine from povidone iodine through either intact skin or broken skin may interfere with thyroid function tests. Contamination with povidone iodine of several types of tests for the detection of occult blood in faeces or blood in urine may produce false-positive result.

Pregnancy and Lactation: Iodine freely crosses the placenta and is secreted in breast milk. Thyroid function disorders have been reported in the offspring of mothers exposed to pharmacological doses of iodine. Povidone iodine should not be used regularly during pregnancy unless there is no alternative treatment available.

Side Effects: Rarely irritation may occur.

Symptoms and Treatment of Overdose: Deliberate or accidental ingestion of large quantities of povidone iodine will result in high blood concentrations of iodine and gastrointestinal corrosive effects including vomiting, diarrhea and abdominal pain. Systemic toxicity may result in shock, hypotension, tachycardia, fever, metabolic acidosis and renal impairment. Symptomatic and supportive treatment should be started with special attention to monitoring electrolyte balance, renal function, thyroid function and liver function. Haemodialysis effectively clears iodine and should be employed in severe cases of iodine poisoning particularly if renal failure is present. Continuous vanovenous haemodiafiltration is less effective than haemodialysis.

Pharmacodynamics: The povidone iodine complex is active at pH values between 2 and 7. The microbial effect is based on the fraction of free, non- complex- bound iodine released in water-based ointments or solutions from the povidone iodine complex as defined in an equilibrium reaction. The povidone iodine complex thus to some degree represents an iodine depot, which release elementary iodine protractedly and in this way guarantees a steady concentration of the efficacious free iodine.

As a result of the binding to the povidone complex, the topical irritant properties of iodine are reduced compared to the alcoholic iodine solutions.

The free iodine reacts as a strong oxidizing agent principally at the molecular level with unsaturated fatty acids as well as readily oxidisable SH or OH groups of the amino acids in enzymes and structural building blocks of microorganism. This non-specific

mode of action explains the comprehensive efficacy of povidone iodine against a wide spectrum of human pathogenic microorganisms, e.g. Gram-positive and Gram-negative bacteria, mycobacteria, fungi (particularly Candida), numerous viruses and some protozoa. Bacterial spores and several species of virus are, in general, inactivated only after longer exposure to an adequate extent.

Specific primary forms of resistance against povidone iodine and also the formation of secondary forms of resistance in long-term application are not anticipated.

Pharmacokinetics: After applying povidone iodine, the possibility of iodine absorption must be considered. This depends upon the nature and duration of treatment as well as the amount applied. Following application to the intact skin only very small amounts of iodine are absorbed. Marked absorption of iodine can occur after long-term application of povidone iodine containing medication to mucosal membranes, extensive skin, wound or burn surfaces and especially after irrigation of body orifices. An elevated iodine concentration in the blood as a result of this is generally transient. In people with a healthy thyroid gland, the increased availability of iodine does not lead to clinically relevant changes in thyroid hormone status. If iodine metabolism is normal, iodine elimination via the kidneys is enhanced.

The absorption of povidone and, to a greater extent, the renal elimination of povidone is dependent on the average molecular weight of the mixture. Above a molecular weight of 35,000 to 50,000 retention within the reticuloendothelial system is to be expected. The accumulation of povidone in the body and other changes which may be seen following intravenous or subcutaneous administration of povidone containing medicaments do not occur after topical application of povidone iodine.

Storage Condition: Store below 30°C. Protect from light. Keep out of reach of children.

Shelf Life: 3 years

Dosage Form: Liquid

Packing size: 15ml, 30ml, 60ml

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

Komposisi: Povidone Iodin 10%w/v

Produk Deskripsi: Larutan berwarna coklat pekat dengan bau tersendiri.

Indikasi: Untuk merawat luka, kulit melecet dan gangguan kulit.

Cara Penggunaan: Sapu di bahagian luka 2 kali sehari.

Laluan Pemberian Ubat: Topikal

Kontraindikasi: Hipersensitiviti kepada povidone iodine.

Amaran dan Perhatian: Penggunaan Povidone Iodin pada kawasan kulit yang rosak harus dielakkan kerana penyerapan iodine berlebihan mungkin berlaku.

Kemerahan atau bengkak yang disebabkan oleh hipersensitiviti, hentikan penggunaan dan rujuk kepada doktor.

Untuk kegunaan luar sahaja.

Interaksi dengan ubat lain: Sila rujuk kepada penerangan Bahasa Inggeris.

Kehamilan dan Penyusuan: Sila rujuk kepada penerangan Bahasa Inggeris.

Kesan Sampingan: Iritasi kulit

Simptom dan Rawatan untuk Dos Berlebihan: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakodinamik: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik: Sila rujuk kepada penerangan Bahasa Inggeris.

Keadaan Penyimpanan: Simpan di bawah 30°C. Lindungi dari cahaya. Jauhkan daripada capaian kanak-kanak.

Tempoh Hayat: 3 tahun

Dos: Cecair

Saiz Bungkus: 15ml, 30ml, 60ml

Maklumat yang terkandung di dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

成份: 优碘10%w/v

外观: 带有独特气味的褐色液体。

主治: 伤口、割伤、擦伤、烫伤及皮肤感染。

用法: 把此药水擦在患处，每天2次。

给药途径: 外用

禁忌: 对优碘过敏者。

警告及注意: 应避免使用优碘在大面积的受损皮肤上，因为可能会出现碘过量吸收。

若使用后因过敏而引起皮肤红肿，请停止使用此药，并咨询医生。

仅限外用。

药物相互作用: 请参阅英文版本。

怀孕和哺乳: 请参阅英文版本。

副作用: 可能会造成皮肤敏感。

药物过量症状及治疗: 请参阅英文版本。

药效学: 请参阅英文版本。

药代动力学: 请参阅英文版本。

贮存: 储藏低于30°C。避免阳光直接照射。远离儿童。

有效期: 3年

剂型: 液体

包装: 15毫升、30毫升、60毫升

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。

Manufactured by Product Registration Holder:

Syarikat Wen Ken Drug Sdn. Bhd.
No. 24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Pengedar Singapura/ Singapore Distributor:

Wen Ken Marketing (S) Pte Ltd
2, Alexandra Road, #02-08, Delta House,
Singapore 159919.

Revision Date: 25-11-2020

...