

건 명	케넨서(Kenhancer) 인서트(말레이시아)		
규 격 (가로x세로)	140x230mm	Q A	음성
		표시문안	6.5pt
색 상	먹 1도	후 가 공	
업체제작일	2023.05.31.	작 업 처	(주)트리스톤

주 요 변 경 내 용	1. 문안 변경, Revised date 변경, 관리번호 변경
-------------------	---------------------------------------



칼선 PANTONE 185C

※ 인쇄진행시, 인쇄사양과 다를경우
녹십자로 즉시 연락주시기 바랍니다.

Analgesics & Anti-inflammatories

Kenhancer^{Plaster}

(Ketoprofen)

[Pharmacological information]

Kenhancer Plaster, targeting on the pain area, has an analgesic and anti-inflammatory effect using TDDS (Transdermal Drug Delivery System) technology. Kenhancer Plaster is characterized in the followings: -
1. Better efficacy : High permeation of Ketoprofen into the pain area
2. Lower incidence of GIT side effects : Avoidance of common side effects(GI) caused by oral Ketoprofen
3. Better absorption: Transcutol® containing in Kenhancer Plaster enhances the permeation of Ketoprofen into skin.

[Composition] 1 sheet (7cmX10cm) contains
Ketoprofen 30mg

[Description]
Plaster which is spread with pale yellowish solid on apricot color of lint fabric.
Each Kenhancer Plaster is a transdermal patch of 7cm X 10cm.

[Indication]
For the treatment of the sign and symptoms of the followings;Arthritis deformans, periarthritis humero-scapularis, tendinitis, peritendinitis, sore muscle, swelling, pain resulted from trauma (contusion, distorsion, sprain etc.)

[Dosage & administration]
Apply 1 sheet of Kenhancer Plaster to affected area twice a day.

[Contraindication]
1. Do not apply to the patients of the following: -
i) Patients with history of hypersensitivity to this drug.
ii) Patients with aspirin induced asthma (nonsteroidal anti-inflammatory drug induced asthma) or with history of it.
2. Carefully apply to the following patients: -
Patients with bronchial asthma.

[Warning and Precautions]
General precautions:
1. Treatment of analgesics & anti-inflammatory drug is not a causal-therapy, but a symptomatic-therapy.
2. In case of using for treatment of inflammation followed skin infection, appropriate antibacterial or antifungal agent should be administered cautiously at the same time, because the effects of ketoprofen may mask the usual symptoms of skin infection.
3. Ketoprofen should be used with caution and under close supervision in case of using this for chronic disease (rheumatoid arthritis, ect). Ketoprofen should also be administered to the patients as the adjunctives with other anti-inflammatory agent.

Precaution to children:
It is not established whether using Ketoprofen for children is safe or not.

Precaution in use:
1. Do not apply to eye and mucous membrane.
2. Temporary pain of irritation if skin may occur when the epidermis is deficient. In such a case, it should be administered cautiously.
3. Should a rash occur after plaster application, treatment should be stopped.
4. Do not expose treated skin to sunlight.

[Pregnancy and Lactation]
As safety in human pregnancy has not established, this preparation should not be administered the wide or excessive use during the prolonged term to pregnant woman or woman of child bearing potential.

[Side effects]
1. Skin: Occasionally redness, rashes, pruritis, dryness and irritation, swelling and photosensitivity may occur. If this symptoms are serious, treatment with this preparation should be discontinued.

[Storage]
Store at below 30°C, protect from moisture and out of direct sunlight.
Keep out of Reach of Children.
Jauhi dari kanak-kanak.

[Packing Size] 1 sheet (7cm X 10 cm) X 6 sheets.

MANUFACTURED BY:
 **GC Biopharma**
GC Biopharma Corp.
26, Mugeuk-ro 65beon-gil, Geumwang-eup,
Eumseong-gun, Chungcheongbuk-do, Korea

PRODUCT REGISTRATION HOLDER:
PHARMA NORD SDN BHD
29-07, Tingkat 29, Q-Sentral,
Jalan Stesen Sentral 2,
Kuala Lumpur Sentral,
50470 Kuala Lumpur, Malaysia

Ins-10-4101-10

Revised date: June 2023