

Three Legs Wintergreen Liniment

Linimen Wintergreen Cap Kaki Tiga



Composition: Methyl Salicylate 25%w/v

Product Description: A clear, slight viscous liquid with smell of methyl salicylate.

Indication: For symptomatic relief of strains, sprains and muscular pains.

Direction for use: Rub on affected area 3 times a day.

Route of Administration: Topical

Contraindications:

Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

This product contains methyl salicylate and when applied or rub on the skin, can be absorbed through the skin into blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Do not use on open wounds. Do not bandage the area after application of this medication.

For external use only. Not to be taken.

Effects on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments:

This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.

Statement on Usage During Pregnancy and Lactation:

Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse Effects/ Undesirable Effects:

May cause slight skin irritation.

Symptoms and Treatment of Overdose:

Mild chronic salicylate intoxication or salicylism usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, swearing, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage.

Pharmacodynamics:

Methyl Salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction.

Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics:

- Methyl Salicylate
- May be absorbed through intact skin.
- The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity.
- Absorption characteristics of salicylates vary with the dose, formulation and route of administration.
- Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin.
- Both the rate and extent of absorption increases after repeated application; increasing the bioavailability.
- Methyl salicylate is extensively metabolized to salicylic acid in the dermal and subcutaneous tissues after topical application.
- At therapeutic levels, the half-life of salicylates is 2-4 hours. As salicylate level increase into the toxic range, the half- life can be greater than 18 hours.

Storage Conditions: Store below 30°C. Keep out of reach of children.

Shelf Life: 3 years

Dosage Form: Liniment

Packing Size: 30ml

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

Komposisi: Metil Salisilat 25%w/v

Deskripsi: Cecair yang jernih, sedikit likat dengan bauan metil salisilat.

Indikasi: Untuk melegakan gejala- gejala akibat seluluh, sengal otot dan sakit otot.

Cara Penggunaan: Gosok pada bahagian terbabit 3 kali sehari.

Laluan Pemberian Ubat: Topikal

Kontraindikasi:

Pesakit yang hypersensitif kepada mana- mana bahan dalam ubat ini.

Amaran dan Perhatian:

Ubat ini mengandungi methyl salicylate dan ia mungkin boleh diserap masuk ke dalam darah melalui kulit. Penyalupan yang berlebihan oleh pesakit yang mengambil warfarin mungkin meningkatkan risiko pendarahan.

Jangan sapu pada luka terbuka. Jangan membalut kawasan luka selepas penggunaan ubat ini.

Untuk kegunaan luar sahaja. Tidak Boleh dimakan.

Kesan ke atas Keupayaan Memandu dan Penggunaan Mesin: Tidak diketahui

Interaksi dengan ubat lain: Sila rujuk kepada penerangan Bahasa Inggeris.

Kehamilan dan Penyusuan: Sila rujuk kepada penerangan Bahasa Inggeris.

Kesan Sampingan: Boleh menyebabkan kerengsaan kulit yang ringan.

Siptom dan Rawatan untuk Dos Berlebihan: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakodinamik: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik: Sila rujuk kepada penerangan Bahasa Inggeris.

Kadaan Penyimpanan: Simpan di bawah 30°C. Jauhkan daripada capaian kanak- kanak.

Tempoh Hayat: 3 tahun

Dos: Linimen

Saiz Bungkus: 30ml

Maklumat yang terkandung di dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

成份: 冬青油 25%w/v

外观: 透明、轻微粘稠及带有水杨酸甲酯气味的液体。

主治: 解除扭伤, 肌肉疼痛和肌肉酸痛。

用法: 搽此药于患处, 一天三次。

给药途径: 外用

禁忌: 对此药成份过敏者不宜使用。

警告及注意:

此产品含有水杨酸甲酯, 当涂抹或擦在皮肤时, 可以通过皮肤渗透进入血液。对于服用华法林的患者, 当因肌肉或关节疼痛时而过度涂抹于皮肤时可能会增加出血的可能性。

不要涂抹于伤口。使用后不要用绷带包扎患处。

外用忌食。

对驾驶和使用机器的能力的影响: 未知

药物相互作用: 请参阅英文版本。

怀孕和哺乳: 请参阅英文版本。

副作用: 可能会引起轻微的皮肤刺激。

药物过量症状及治疗: 请参阅英文版本。

药效学: 请参阅英文版本。

药代动力学: 请参阅英文版本。

贮存: 储藏低于30°C。远离儿童。

有效期: 3年

剂型: 搽剂

包装: 30毫升

此说明书仅属概述, 欲知更详细资料, 可咨询您的药剂师或医生。

Manufactured by Product Registration Holder:

Syarikat Wen Ken Drug Sdn.Bhd.
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Singapore Distributor:

Wen Ken Marketing (S) Pte. Ltd.
2, Alexandra Road, #02-08 Delta House,
Singapore 159919.

Date of Revision: 03.12.2020