Filename : MY-BRU_Yellow-Balsem_Leaflet_27082025
Client : SWKD
Requestor : -
Pdt. PIC : Shu Ying
Designer : Nickles
Received : 06.05.2025
Amend : 27.08.2025
Version : 01
Date FA : -
Opened : 127w x 160h MM
Closed : -
Colour : 1C x 1C
K

THREE LEGS YELLOW BALSEM
BALSEM KUNING CAP KAKI TIGA
三脚标青草膏

THREE LEGS YELLOW BALSEM

Composition: Methyl Salicylate 13.5%w/w, Menthol 7.5%w/w, Camphor 7%w/w

Product Description: A yellow colour ointment with menthol smell.

Indication: For symptomatic relief of muscular pain, rheumatic pain and sprain.

Direction for use: Rub on affected area 2 to 3 times a day.

Route of Administration: Topical

Contraindications: Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions: This product can cause convulsion. This product is contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor-containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

This product contains methyl salicylate and when applied or rub on the skin, can be absorbed through the skin into blood.

For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

If you are taking warfarin, please consult your doctor before using this product.

Avoid contact with the eyes and mucous membrane. Do not apply to open wounds or damaged skin. Do not apply over large areas of the body or bandage or use with heating pads.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.

For external use only. Not to be taken.

Effects on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments: This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.

Statement on Usage During Pregnancy and Lactation: Do not use during pregnancy or when breastfeeding

Adverse Effects/ Undesirable Effects: Skin irritation, redness or swelling, burning or stinging sensation. Adverse reaction is rare, symptom of dizziness, nausea may occur.

Symptoms and Treatment of Overdose: Mild chronic salicylate intoxication or salicylism usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, swearing, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage.

Pharmacodynamics: Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol and methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate is a salicylic acid derivative.

Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction.

Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics:

Camphor

- Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure.
- It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine.
- Camphor crosses the placenta.

Menthol

- After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate

- May be absorbed through intact skin.
- The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity.
- Absorption characteristics of salicylates vary with the dose, formulation and route of administration.
- Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin.
- Both the rate and extent of absorption increases after repeated application; increasing the bioavailability.
- Methyl salicylate is extensively metabolized to salicylic acid in the dermal and subcutaneous tissues after topical application.
- At therapeutic levels, the half-life of salicylates is 2-4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Storage Conditions: Store below 30°C. Protect from light. Keep out of reach of children.

Shelf Life: Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Ointment

Packing Size: 20g, 36g

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

BALSEM KUNING CAP KAKI TIGA

Komposisi: Metil Salisilat 13.5%w/w, Mentol 7.5%w/w, Camphor 7%w/w

Deskripsi: Salap yang berwarna kuning dan berbau menthol.

Indikasi: Untuk melegakan sakit otot, sakit reumatik dan sakit urat.

Cara Penggunaan: Sapu pada bahagian terbabit 2 hingga 3 kali sehari.

Laluan Pemberian Ubat: Topikal

THREE LEGS YELLOW BALSEM
BALSEM KUNING CAP KAKI TIGA
三脚标青草膏

Kontraindikasi: Pesakit yang hypersensitif kepada mana-mana bahan dalam ubat ini.

Amaran dan Perhatian: Ubat ini boleh menyebabkan sawan dan tidak sesuai untuk bayi di bawah umur 2 tahun dan perhatian perlu diberikan kepada kanak-kanak berumur 2 tahun ke atas apabila menggunakannya. Sapu ubat yang mengandungi camphor pada hidung kanak-kanak adalah merbahaya kerana boleh menyebabkan pengsan.

Ubat ini mengandungi methyl salicylate dan ia mungkin boleh diserap masuk ke dalam darah melalui kulit.

Penyapuan yang berlebihan oleh pesakit yang mengambil warfarin mungkin meningkatkan risiko pendarahan.

Sekiranya anda sedang mengambil warfarin, sila rujuk kepada doktor anda sebelum menggunakan produk ini.

Tidak boleh sapu pada bahagian mata, permukaan mukus, luka terbuka atau kulit yang rosak. Elakkan penggunaan yang berlebihan secara meluas pada bahagian badan atau membalut.

Pada mulanya sapu kuantiti yang kecil pada kawasan kulit yang terhad dan memerhati sama ada ketidakselesaan atau gatal berlaku sebelum sapu pada kawasan kulit yang lebih besar. Jika iritasi kulit berlaku, sila berhenti gunakannya.

Untuk kegunaan luar sahaja. Tidak boleh dimakan.

Kesan ke atas Keupayaan Memandu dan Penggunaan Mesin: Tidak diketahui

Interaksi dengan ubat lain: Sila rujuk kepada penerangan Bahasa Inggeris.

Kehamilan dan Penyusuan: Sila rujuk kepada penerangan Bahasa Inggeris.

Kesan Sampingan: Iritasi kulit, kemerahan, bengkak, rasa pedih berlaku. Reaksi buruk jarang berlaku, simptom pening, loya boleh berlaku.

Simptom dan Rawatan untuk Kelebihan Dos: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakodinamik: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik: Sila rujuk kepada penerangan Bahasa Inggeris.

Keadaan Penyimpanan: Simpan di bawah 30°C. Lindungi daripada cahaya. Jauhkan daripada capaian kanak-kanak.

Tempoh Hayat: Sila rujuk pada pembungkusan. Jangan guna selepas tarikh luput.

Dos: Salap

Saiz Bungkusan: 20g, 36g

Maklumat terkandung dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

三脚标青草膏

成份： 冬青油 13.5%w/w，薄荷 7.5%w/w，樟脑 7%w/w

外观： 带有薄荷气味的黄色软膏。

主治： 解除肌肉疼痛，风湿疼痛及扭伤。

用法： 擦在患处，每天2至3次。

给药途径： 外用

禁忌： 对此药成份过敏者不宜使用。

警告及注意： 本产品含有樟脑，可能会引起抽搐，两岁以下婴儿不宜使用。用于较年长儿童时需格外谨慎。将含有樟脑的产品涂抹于儿童的鼻孔是非常危险的，即使少量使用也可能导致休克。

此产品含有水杨酸甲酯，当涂抹或擦在皮肤时，可以通过皮肤渗透进入血液。

对于服用华法林的患者， 当因肌肉或关节疼痛时而过度涂抹于皮肤时可能会增加出血的可能性。

如果您服用华法林，请在使用本产品前咨询您的医生。

避免与眼睛和粘膜接触。不要涂抹于伤口或受损皮肤。不要大面积涂抹在身体或用绷带包扎或使用加热垫。

在涂抹于较大区域的皮肤前，涂抹少量在有限的皮肤区域以观察是否有不适或瘙痒。若刺激反应持续，请停止使用。

外用忌食。

对驾驶和使用机器的影响： 未知

药物相互作用： 请参阅英文版本。

怀孕和哺乳： 请参阅英文版本。

副作用： 可能会造成皮肤敏感。

药物过量症状及治疗： 请参阅英文版本。

药效学： 请参阅英文版本。

药代动力学： 请参阅英文版本。

贮存： 储藏低于30°C。避免阳光直接照射。远离儿童。

有效期： 请参阅包装。请勿在有效期后使用。

剂型： 软膏

包装： 20毫克、36毫克

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。

Date of Revision: AUGUST 2025

Manufactured by Product Registration Holder:
Syarikat Wen Ken Drug Sdn. Bhd.
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Singapore Distributor:
Wen Ken Marketing (S) Pte. Ltd.
2, Alexandra Road, #02-08 Delta House,
Singapore 159919.