

Skilax Drops 7.5mg/ml

Sodium Picosulfate

Composition per ml:

Sodium Picosulfate 7.5 mg

(each ml is approx. equiv. to 20 drops)

Pharmaceutical form and content:

Solution for oral use in drops. Bottle of 15 ml.

Product Description:

Clear, colourless to slightly yellowish odourless liquid.

Pharmacodynamics:

Sodium Picosulfate has a local laxative effect, which will stimulate the mucosa of both the large intestine and the rectum after bacterial cleavage in the colon. This results in colonic peristalsis by promoting accumulation of water and electrolytes in the colonic lumen. This stimulation and osmotic effect cause defecation, reduction of transit time and softening of the stool. Stimulation of the rectum causes increased motility and a feeling of rectal fullness.

Sodium Picosulfate has no effect on alteration of digestion or absorption of calories and essential nutrients in the small intestine.

Pharmacokinetics:

Absorption and Distribution

After oral absorption, Sodium Picosulfate reaches the colon with no significant absorption and systemic effect.

Biotransformation

Sodium Picosulfate is converted into the active form, bis-(p-hydroxyphenyl)-pyridyl-2-methane (BHPM), via bacterial cleavage in the distal segment of the intestine. The onset of action of the preparation is usually between 6 - 12 hours. The laxative effect is independent of the plasma levels of the active moiety.

Elimination

Only small amounts of the active form are absorbed and almost completely conjugated in the intestinal wall and liver to form the inactive BHPM glucuronide. After oral administration of 10mg Sodium Picosulfate 10.4% of the total dose was excreted as BHPM glucuronide in urine after 48 hours. In general, urinary excretion is inversely proportional to the concentration of Sodium Picosulfate administered.

Indication:

For the management and short term relief of constipation of any aetiology.

Recommended Dose:

Adults and children over 10 years old: 5 - 10mg (14 to 27 drops per day).

Children under 10 years old: Not to be taken by children under 10 years old without medical advice.

Children 4 to 10 years old: 2.5 – 5mg (7 to 14 drops per day).

Children under 4 years old: The recommended dosage is 0.25mg per kilogram body weight per day (1 ml of Skilax Drops 7.5mg/ml contains 7.5mg Sodium Picosulfate).

Doses should preferably be taken before going to bed. The effect will be noticed next day in the morning. (Each ml is approx. to 20 drops)

Contraindications:

Skilax is contraindicated in patients with:

- Ileus or intestinal obstruction
- Severe painful and/or feverish acute abdominal conditions (e.g. appendicitis) potentially associated with nausea and vomiting
- Acute inflammatory bowel diseases
- Severe dehydration
- Known hypersensitivity to Sodium Picosulfate or any other component of the product
- Rare hereditary conditions that may be incompatible with an excipient of the product.

Warnings & Precautions:

Skilax should not be taken continuously for more than five days without knowing the cause of constipation. Prolonged, excessive use may lead to fluid and electrolyte imbalance and hypokalaemia.

This product should not be taken by children under 10 years old without medical supervision.

Interaction with Other Medicaments:

The concomitant use of diuretics or adreno-corticosteroids may increase the risk of electrolyte imbalance if excessive doses of Skilax are taken.

Electrolyte imbalance may lead to increased sensitivity to cardiac glycosides.

Concurrent administration of antibiotics may reduce the laxative action of this product.

Use during Pregnancy and Lactation:Pregnancy

There are no adequate and well-controlled studies in pregnant women. Long experience has shown no evidence of undesirable or damaging effects during pregnancy.

Lactation

Clinical data show that neither the active moiety of sodium picosulfate (BHPM or bis-(p-hydroxyphenyl)-pyridyl-2-methane) nor its glucuronides are excreted into the milk of healthy lactating females. Nevertheless, as with all medicines, Skilax should not be taken during pregnancy, especially in the first trimester, and during breast feeding unless the expected benefit is thought to outweigh any possible risk and only on medical advice.

Side Effects:

Side effects include diarrhea, abdominal discomfort, abdominal cramps, nausea, vomiting and dizziness.

Skilax may also cause hypersensitivity and skin reactions such as angioedema, drug eruption, rash and pruritus.

Symptoms and Treatment of Overdose:

Overdose would lead to abdominal cramps and profuse diarrhea and loss of fluid and electrolytes. Chronic abuse may lead to secondary hyperaldosteronism, renal calculi and hypokalaemia with associated renal tubular damage, metabolic alkalosis and muscle weakness. Isolated cases of colonic mucosal ischaemia have been observed with overdoses of Sodium Picosulfate.

Treatment includes immediate induction of vomiting or by gastric lavage to minimize drug absorption. Replacement of fluids and correction of electrolyte imbalance may be required especially in the elderly and the young. Antispasmodic agents may be used if necessary.

Storage Conditions: Store below 30°C. Protect from light.

Packaging: 15ml/ bottle

Registration No.: MAL20016154XC

Product Registration Holder :
Milrin Pharmaceutical Co. (M) Sdn. Bhd.
198101009339 (75454-P)
125L, 5th Floor, Wisma Getah,
Macalister Road, 10400 Georgetown,
Penang, Malaysia.

Manufacturer:
TERAPUTICS SDN. BHD.
(590500-W)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS, Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Date of Revision: 13 January 2020
MYX026PI(01)