

Vimelinspec

Solution for injection

COMPOSITION:

Each ml contains:
Lincomycin (as HCl)50mg
Spectinomycin (as sulfate)100mg
With 1.0% (v/v) BenzylAlcohol as a preservative

DESCRIPTION

The finished product is colorless to yellowish transparent solution

PHARMACODYNAMICS

The lincomycin-spectinomycin combination covers a broad spectrum of activity against gram-positive, gram-negative and mycoplasma bacteria. Lincomycin is especially active against gram-positive bacteria (such as *Streptococci* and *Staphylococci*) and mycoplasmas, while spectinomycin is more specifically active against gram-negative bacteria (*E. coli*, *Pasteurella*, *Salmonella*), mycoplasmas and gram-positive bacteria. Resistance to lincomycin is often conferred by plasmid factors (erm-genes) encoding ribosomal binding site-modifying methylases and frequently cross-resistance to other MLS_B-group antimicrobials. However, the most common mechanism in mycoplasmas is alteration of the binding site by mutational events (chromosomal resistance). Resistance to lincomycin, mediated by efflux pumps, or inactivation of enzymes, has also been described. There is often complete cross-resistance between lincomycin and clindamycin. High level resistance to spectinomycin develops by one-step chromosomal mutation in many enteric bacteria (such as *E. coli*). Plasmid resistance is less common. Strains with chromosomal resistance do show not cross-resistance with aminoglycosides.

PHARMACOKINETICS

Lincomycin is well distributed throughout the body and is metabolized significantly. Spectinomycin is also well distributed throughout the body and appears to be mainly excreted as the parent compound.

INDICATION

Treatment of infections caused by germs sensitive to lincomycin and / or spectinomycin.

In calves: Treatment of respiratory infections, arthritis and omphalitis.

In sheep and goats: Treatment of respiratory infections and mycoplasmosis.

In pigs:

- Treatment of intestinal adenomatosis (ileitis) due to

Lawsonia intracellularis,

- Treatment of haemorrhagic enteritis and colibacillosis,

- Treatment of mycoplasmosis,

- Treatment of infectious arthritis.

In chickens and turkeys:

- Treatment of mycoplasmosis associated or not with

Escherichia coli,

- Treatment of aerococculitis due to *Escherichia coli*,

- Treatment of avian cholera due to *Pasteurella multocida*.

In cats and dogs:

- Treatment of respiratory, intestinal, urinary infections, skin infections (including wounds and abscesses) and arthritis, caused by germs susceptible to lincomycin and / or spectinomycin, including *Staphylococcus* spp., *Streptococcus* spp., *Bacteroides* spp., *Clostridium* spp., *Fusobacterium* spp., *Actinomyces* spp., *Mycoplasma* spp.

TARGET SPECIES

Cattle (calves), sheep, goats, pigs, chickens, turkeys, cats and dogs.

ROUTE OF ADMINISTRATION

By intramuscular or subcutaneous injection

RECOMMENDED DOSE / USE INSTRUCTION

The product should be administered by intramuscular or subcutaneous injection. In order to ensure a correct dosage, the weight of the animal should be determined as accurately as possible.

Calve : 5 mg lincomycin and 10 mg spectinomycin per kg body weight (corresponding to 1 ml of solution per 10 kg body weight) intramuscularly, two injections during the first day of treatment, then one injection per day for 3 to 5 days.

Sheep and goats: 5 mg lincomycin and 10 mg spectinomycin per kg body weight (corresponding to 1 ml solution per 10 kg body weight) intramuscularly, once daily, for 3 to 5 days.

Pigs: 5 mg lincomycin and 10 mg spectinomycin per kg bodyweight (corresponding to 1 ml solution per 10 kg body weight) intramuscularly, repeat, if necessary, 24 hours later and for a maximum of 5 days.

Chickens and turkeys: 10 mg of lincomycin and 20 mg of spectinomycin per kg bodyweight (corresponding to 1 ml of solution per 5 kg of weight) subcutaneously, once a day for 3 days.

Cats and dogs: 10 mg lincomycin and 20 mg

spectinomycin per kg body weight (corresponding to 1 ml solution per 5 kg body weight intramuscularly) Treatment may be repeated between 12 and 24 hours apart for 3 at 7 days depending on the clinical response.

CONTRAINDICATION

Do not use in case of hypersensitivity to the active substances or to any of the excipients. Do not use in rabbits, hamsters, guinea pigs, chinchillas or horses, as this may result in severe gastrointestinal disturbances.

WARNINGS AND PRECAUTIONS

i) Special precautions for use in animals The veterinary medicinal product must be used on the basis of susceptibility tests of bacteria isolated in animals. If this is not possible, the therapy should be based on local epidemiological information (region, farm level) regarding the sensitivity of the target bacteria.

Use of the product as per the recommendations may increase the prevalence of resistant bacteria and may decrease the effectiveness of treatment with macrolides given the potential risk of cross-resistance.

Use with caution in animals that have already shown signs of allergy.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Lincomycin and spectinomycin may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact.

Allergic reactions to these substances can be serious. Precautions must be taken to avoid accidental self-injection.

In case of accidental self-injection, seek medical advice immediately and show the label or package leaflet.

Avoid all contact with skin and eyes. Wash off spills immediately with plenty of water. Wash hands after use.

People with known hypersensitivity to these antibiotics should avoid contact with the veterinary medicinal product.

If symptoms occur after exposure to the product, such as a rash, seek medical advice immediately and show the package leaflet or the label to the physician. Swelling of the face, lips or eyelids, or breathing difficulties are more serious symptoms and require urgent medical attention.

iii) Other precautions

Any.

- Keep out of reach of children
- Jauhkan daripada kanak-kanak

INTERACTIONS WITH OTHER MEDICAMENTS

Do not administer with macrolides.

Combination with anesthetics may lead to neuromuscular blockade.

PREGNANCY AND LACTATION

The safety of the product has not been established in pregnant females, lactating or laying eggs. The use of this product should be done only after evaluation of the benefit / risk ratio, established by the responsible veterinarian.

SIDE EFFECTS / ADVERSE REACTIONS

A softening of the stool can be observed. It is usually transient and retrocedes in a few days without treatment.

Use of this product may cause occasional discomfort during injection.

SIGNS AND SYMPTOMS OF OVERDOSE AND TREATMENT

Not known

STORAGE CONDITIONS

Store in a cool dry place not exceeding 30°C. Protect from light.

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 24 hours

WITHDRAWAL PERIOD

Meat and offal: 14 days (Calves, sheep, goats, pigs, chickens and turkeys)

Milk: Do not use in females producing milk for human consumption during lactation or dry period, or for future producers of human milk during the 2 months prior to farrowing.

Eggs: do not use in laying birds of eggs for human consumption.

PACKAGING: 100ml

REVISED DATE: Dec 04, 2019