

[말레이시아]
하이알프리필드주 설명서

Size : 130 × 200 (mm)

20/07/29 수정사항

- 1. HOW TO USE 항목 추가 및 이미지 삽입
- 2. 설명서 사이즈 변경
(120x175 → 130x200(mm))

21/01/29 수정사항

- 1. ROUTE OF ADMINISTRATION 추가
- 2. ADVERSE REACTIONS→ ADVERSE EFFECTS/
UNDESIRABLE EFFECTS 변경
- 3. PRECAUTIONS -> WARNING AND PRECAUTIONS
- 4. INTERACTION WITH OTHER MEDICAMENTS 추가
- 5. OVER DOSE AND TREATMENT 추가
- 6. INCOMPATIBILITIES 추가

22/10/21 수정사항

- 1. HOW TO USE 오기건 수정
- 2. Date of Revision 추가

For the Improvement of Joint Disease

HYAL Prefilled **L** Injection
Sodium Hyaluronate

HYAL Injection composed of sodium hyaluronate that is available for the intra-articular injective treatment of osteoarthritis of the knee and periarthritis of shoulder relieves pain and has metabolic effects on articular cartilage, synovial tissue, and synovial fluid. Additionally, with prefilled syringe system, HYAL Injection is a safe and effective treatment for patients.

PHARMACOLOGY

At six and 12 weeks after the initiation of administration, the test group (administered sodium hyaluronate) was compared with a group administered physiological saline solution to determine the effects on articular cartilage and synovial tissue.

In both the six and 12-week period, cartilage degeneration proceeded with the lapse of time in both groups: however at 12 weeks, the efficacy of HYAL Inj. (sodium hyaluronate) in inhibiting degeneration was clearly observed at the lateral condyle of the femur and tibia where relatively marked degeneration was observed in the saline group. The effects of the intra-articular sodium hyaluronate(HA) injection on the osteoarthritis knee joint was evaluated in dogs and using an experimental model of osteoarthritis induced by sectioning the anterior cruciate ligament. Seven weeks after surgery, the damage was significantly reduced in knee joints treated with HA from the second week postsurgery compared to untreated joints.

PHARMACOKINETICS

Fluorescence distribution was investigated in two models of experimentally-induced osteoarthritis in rabbits after intra-articular administration of fluorescein-labeled sodium hyaluronate, osteoarthritic cartilage showed fluorescence distributed in the deeper layers of cartilage, which presented marked fibrillation, cleft formation, and loss of staining with safranin O. Synovial tissue in osteoarthritis induced by surgical dissection of the anterior cruciate ligament showed hyperplasia and hypertrophy, and inflammatory cell infiltration was observed. These results suggest that fluorescein-labeled sodium hyaluronate administered into the joint cavity can penetrate through the osteoarthritis cartilage. In the phase 1 study, high molecular weight sodium hyaluronate(HA) was administered intra-articularly at a single dose of 15-50mg in the knee in 29 healthy male volunteers and its pharmacokinetics evaluated. The serum HA concentration changed in the range of about 30-150ng/mL after administration of HA at each dose or the placebo. Thus, the serum HA concentration did not significantly exceed its physiological range after administration of exogenous HA into the knee.

■ **DESCRIPTION**

HYAL Injection comes in a 2.5mL pre-filled syringe. It is colorless, clear and viscous. Each pre-filled syringe comes in a printed box with a package insert.

■ **COMPOSITION**

Each ml contains 10mg of sodium hyaluronate

■ **INDICATIONS**

Osteoarthritis of knee, Periarthritis of shoulder

■ **DOSAGE AND ADMINISTRATION**

For adults, administer 1 syringe per once to the articular cavity of knee or articular cavity of shoulder, intra-subacromial bursa, intra-biceps tendon sheath five times continuously in weekly intervals. The number of administration times may be adjusted according to symptoms. Administration must be conducted in completely aseptic conditions for intra-articular injection.

■ **ROUTE OF ADMINISTRATION** : Intra-articular


■ **HOW TO USE**



1. Take off the rubber cap of syringe.



2. Put in the needle.



3. Inject after removing the air inside the syringe.



4. Throw away after disconnecting the needle from the syringe.

■ **CONTRAINDICATION**

Patients with a history of hypersensitivity to this drug.

■ **CAREFUL ADMINISTRATION**

- 1) Patients with a history of hypersensitivity to other drugs.
- 2) Patients who are or with a history of hepatic disorder.

■ **ADVERSE EFFECTS/ UNDESIRABLE EFFECTS**

When administered to the knee, pain and inflammation may occur at the injection site. There have also been occasional reports of hypersensitivity, including rarely anaphylaxis.

■ **WARNING AND PRECAUTIONS**

- 1) Since the administration of this drug may cause a local inflammation when the inflammation due to osteoarthritis of the knee is severe, it is desirable to administer this drug after the elimination of inflammation.
- 2) Since local pains may appear, it is recommended to indicate managements such as a local stabilization following to the administration of this drug.
- 3) Since the leakage of this drug to extra-articular cavity may induce pains, administer this drug only to intra-articular cavity.

■ **INTERACTION WITH OTHER MEDICAMENTS**

The safety and effectiveness of this drug and other intra-articular medications have not been proven to be co-administered.

■ **USE DURING PREGNANCY OR LACTATION**

- 1) Teratogenic activities were not reported in animal experiments but the safety in pregnant women is not established. This drug is administered carefully to pregnant or women of child bearing potential.
- 2) As it has been reported in animal studies that this drug is excreted into breast milk, lactation should be avoided during the period of treatment.

■ **USE IN CHILDREN**

Since the safety in children is not established, it is administered carefully when unavoidable.

■ **CAUTION FOR USE**

- 1) Because this drug is administered to the articular cavity of knee or shoulder, administration must be conducted in completely aseptic conditions.
- 2) If no improvement of symptoms appears after administration of 5 times, the administration should be discontinued.
- 3) When articular fluid is stagnated, eliminate articular fluid by use of prickle as necessary.
- 4) It should not be administered to blood vessel.
- 5) It should not be used for ophthalmic remedy.
- 6) It is recommended to use the needle size of 18-20G because this drug is viscous.
(※ In case of absorbing this drug from ampoule.)
- 7) Since this drug may be precipitated by chlorhexidine or the quaternary ammonium salts such as benzalkonium chloride, cautions should be taken.

* **CAUTIONS** *

- 1) Do not use if the rubber cap of syringe has been tampered.
- 2) For single use only. To be discarded once used.

■ **OVER DOSE AND TREATMENT** : Non reported

■ **INCOMPATIBILITIES** : Non reported

■ **SHELF LIFE** : 3 years from manufacturing date

■ **STORAGE** : Store in a tight and light-resistant container at room temperature.
Store below 30°C
Keep medicines out of reach of children.
Jauhkan ubat-ubatan dari kanak-kanak.

■ **HOW SUPPLIED** : 2.5mL(1 Syringe) × 1 Syringe, 5 Syringe

Product Registration Holder :

The Zylas Medical Co: No. 7, Jalan Molek 3/10, Taman Molek, 81100 Johor Bahru

Date of Revision : OCT 2022

