

DOMIDIN®

LOZENGES 0.5MG

Ingredient(s):

Each lozenge contains:

Domiphen bromide 0.5mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

1. Domiphen bromide belongs to the category of long-chained quaternary ammonium compounds. It has a disinfecting effect against most bacteria and fungi that are possible causes of infection of the mouth and pharynx. Among these, the gram-positive microorganisms are more sensitive to Domiphen bromide than the gram-negative ones.
2. In aqueous solutions, Domiphen bromide dissociates into a relatively large and complex cation which is responsible for the surface activity, and a smaller inactive anion. In addition to its emulsifying and detergent properties, Domiphen bromide has bactericidal activity against gram-positive and at higher concentrations, against some gram-negative bacteria in the mouth and throat.
3. The amount of Domiphen bromide that reaches the body fluids after oral administration cannot be measured with normal analytical methods. Current knowledge suggests that the systemic absorption of Domiphen bromide is very low.

Indication(s):

For the relief of sore throat caused by excessive expectoration, stomatitis and pharyngolaryngitis.

Dosage and Administration:

One lozenge to be sucked every 1 to 2 hours, as necessary. Maximum 8 lozenges a day.

Route of administration: Oral

Contraindication(s):

Contraindicated in patients who have shown hypersensitivity to this product or related quaternary ammonium compounds.

Precaution(s) / Warning(s):

1. If the patient's symptoms persist after 5 days of treatment, discontinue the use of the product and immediately contact a doctor or pharmacist for professional assistance.
2. Safe use of this product during pregnancy and lactation has not been established. Women who suspect pregnancy, pregnant women or nursing mothers should not be given this product unless clearly needed.

Drug Interaction(s):

This product is incompatible with anionic surfactants (e.g. soap and tooth paste) and protein containing organic (e.g. blood and pus) which can decrease the disinfecting activity of Domiphen bromide.

Side Effect(s) / Adverse Reaction(s):

Side effects rarely occur consisting of local irritation or reddening of mucous membrane, nausea and vomiting.

Domiphen bromide has depolarizing muscle relaxant properties and toxic symptoms include dyspnoea and cyanosis due to paralysis of the respiratory muscles, possibly leading to asphyxia.

Symptoms and Treatment for Overdosage, and Antidote(s):

Treatment of poisoning is symptomatic, demulcents may be given if necessary.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A white to slight yellowish round tablet, each side with a mark of



Blister packing of 10’s x 10 and 10’s x 100.

LOZENG
DOMIDIN® 0.5MG

Bahan Aktif:

Setiap Lozeng mengandungi:
Domiphen bromide 0.5mg

Indikasi:

Untuk melegakan sakit kerongkong yang disebabkan oleh ekspektorasi berlebihan, stomatitis dan faringolaringitis.

Dos:

Satu lozeng dihisap setiap 1 hingga 2 jam, bila perlu. Maksimum 8 lozeng sehari.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679