



AXCEL[®]

FUSIDIC ACID

OINTMENT

COMPOSITION:

Sodium Fusidate (micronised) 2% w/w

PRESENTATION:

Translucent, waxy, water immiscible ointment.

INDICATIONS:

Skin infection caused by Staphylococci, Streptococci, and other susceptible or sodium fusidate-sensitive organisms.

PHARMACOLOGY:*Mechanism of Action:*

Sodium fusidate is a sodium salt for fusidic acid, an antimicrobial substance produced by the growth of certain strain of *Fusidium coccineum*. Fusidic acid is steroidal antibiotic with a bacteriostatic or bactericidal activity mainly against Gram-positive bacteria. Fusidic acid inhibits bacterial protein synthesis by inhibiting a factor necessary for the translocation of peptide subunits and elongation of the peptide chain. Fusidic acid is very active against staphylococci. Notably *Staph. aureus* and *Staph. epidermidis*, *Nocardia asteroides* and many clostridial strains are also highly susceptible. Most gram-negative bacteria are intrinsically resistant but fusidic acid is active against *Neisseria spp.* *Bacteroides fragilis* is susceptible and it has some activity against strains of *Mycobacterium tuberculosis* and *M. leprae*. Fusidic acid has some activity against a range of protozoa including *Giardia lamblia* and *Plasmodium falciparum*. The therapeutic efficacy of topically applied Sodium Fusidate Ointment is due partly to its pronounced antibacterial activity against the organisms responsible for skin infections, and partly to its ability to penetrate intact skin.

Pharmacokinetics:

Fusidic acid penetrates normal or inflamed skin and achieves bactericidal concentrations down to the lower corium.

DIRECTION FOR USE:

For external use only.

Apply a thin layer of the ointment to the affected area 2 - 3 times a day, generally for a period of seven days.

CONTRAINDICATIONS:

Patients known to be hypersensitivity to Sodium Fusidate. Caution should be taken when applying Axcel Fusidic Acid Ointment in the eye region as this preparation may cause irritation if it gets into

the eye.

PRECAUTIONS:

Repeated topical use of the ointment may lead to problems of resistance.

Use in Pregnancy and Lactation:

The safety of the use during pregnancy and lactation has not been fully established. Therefore, they should not be used extensively on pregnant and nursing mothers, in large amounts or for prolonged periods of time.

Use in Infants and Children:

Avoid prolonged use in infants and children.

SIDE EFFECTS:

Hypersensitivity reactions in the form of rashes and irritation may occur after the topical administration of the ointment. It should be avoided in patients known to be hypersensitive to sodium fusidate or its derivatives.

DRUG INTERACTIONS:

No known drug interaction to date when applied topically.

OVERDOSAGE AND TREATMENTS:

Overdosage is unlikely to occur when sodium fusidate is applied topically.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK****PACK QUANTITIES:**

Available in aluminium tube of 5gm and 15gm.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL20021410AZ



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

REGN - 220817