

Kipo for IV Infusion 200mg/mL

COMPOSITION

Each mL contains: N(2)-L-alanyl-L-glutamine 200mg
(equivalent to L-alanine 82mg, L-glutamine 134.6 mg)
Excipients: Water for Injection

CHARACTERISTICS

Osmolarity	pH	Titration acidity
830 - 1010 mOsm/L	5.4 - 6.0	90.0 to 105.5 mmol NaOH/L

PRODUCT DESCRIPTION

Kipo for IV Infusion 200 mg/ml is a sterile clear and colorless solution, packed in a 50 mL and 100 mL type I clear glass vial and seal with 20mm rubber stopper and 20mm aluminium/plastic flip-off sealcap.

INDICATION

Part of an intravenous parenteral nutrition regimen as a supplement to amino acid solutions or an amino acid containing Infusion regimen, e.g. In patients in hypercatabolic. and/or hypermetabolic states.

PHARMACODYNAMICS

N(2)-L-alanyl-L-glutamine is endogenously split into the amino acids glutamine and alanine, hence supplying glutamine from intravenous nutrition. The released amino acids enter the respective nutrient storage in the body, and are metabolized according to the needs of organisms.

PHARMACOKINETICS

N(2)-L-alanyl-L-glutamine, is split into the amino acids glutamine and alanine rapidly after infusion. The half-life is 2.4 - 3.8 minutes in human (4.2 minutes in end-stage renal failure). The plasma clearance is 1.6 - 2.7 L/min. The corresponding free amino acids increase in equimolar amount as the product disappears. It's probably that hydrolysis takes place completely in extracellular space. Under continuous infusion, the renal elimination of N(2)-L-alanyl-L-glutamine is below 5%, which is similar to other amino acid infusions.

Many patients requiring intravenous nutrition are depleted in glutamine. Therefore, infusion of glutamine supplement is required.

CONTRAINDICATIONS

Do not apply this product to the following patients:

- Severe kidney impairment (clearance of creatinine < 25mL/min)
- Severe liver impairment
- Severe metabolic acidosis
- Patients who are hypersensitive to any ingredient of this product

DOSAGE AND ADMINISTRATION

Route of administration: Intravenous (parental). Administration of this product should only be given by the physicians.

Solution for infusion after mixing with a compatible solution and administered by intravenous infusion. Solutions of mixtures with an osmolarity above 800 mosmol/L should be infused by the central venous route.

Adults:

Kipo for IV Infusion 200mg/mL is administered parallel with parenteral nutrition or enteral nutrition or a combination of both. Dosage depends on the severity of the catabolic state and on amino acids/protein requirement. A maximum daily dosage of 2 g amino acids/or protein per kg bodyweight should not be exceeded. The supply of alanine and glutamine should be taken into consideration in the calculation. The proportion of the amino acids supplied through Kipo for IV Infusion 200mg/mL should not exceed approx. 30% of the total amino acids/protein supply.

Daily dose:

1.5 - 2.5 mL of Kipo for IV Infusion per kg body weight (equivalent to 0.3 – 0.5 g N(2)-L-alanyl-L-glutamine per kg body weight). This equates to 100 to 175 mL Kipo for IV Infusion for a patient of 70 kg body weight.

Maximum daily dose:

2.5 mL equivalent to 0.5 g N(2)-L-alanyl-L-glutamine of Kipo for IV Infusion per kg body weight.

The maximum daily dose of 0.5 g N(2)-L-alanyl-L-glutamine per kg body weight should be administered in combination with at least 1.0 g

amino acids/protein per kg body weight per day. This results in a daily dosage of at least 1.5 g amino acids/protein per kg body weight.

The following adjustments are examples for the supply with Kipo for IV Infusion 200mg/mL and amino acids through the parenteral nutrition solution, and/or protein through enteral nutrition formula:

- *Amino acids/protein requirement 1.2 g/kg body weight per day:*
0.8 g amino acids/protein + 0.4 g N(2)-L-glutamine per kg body weight
- *Amino acids/protein requirement 1.5 g/kg body weight per day:*
1.0g amino acids/protein + 0.5 g N(2)-L-alanyl-L-glutamine per kg body weight
- *Amino acids/protein requirement 2 g/kg body weight per day:*
1.5 g amino acids/protein + 0.5 g N(2)-L-alanyl-L-glutamine per kg body weight

Kipo for IV Infusion 200mg/mL is an infusion solution concentrate which is not designed for direct administration.

Patients with total parenteral nutrition

The rate of infusion depends on that of the carrier solution and should not exceed 0.1g amino acids/kg body weight per hour. Kipo for IV Infusion should be mixed with a compatible amino acid carrier solution or an amino acid containing infusion regimen prior to administration.

Patients with total enteral nutrition

Kipo for IV Infusion is continuously infused over 20-24 hours per day. For peripheral venous infusion, dilute Kipo for IV Infusion to an osmolarity ≤ 800 mosmol/l (e.g. 100 ml Kipo for IV Infusion + 100 ml saline).

Patients with combined enteral and parenteral nutrition

The full daily dosage of Kipo for IV Infusion should be administered with the parenteral nutrition, i.e. mixed with a compatible amino acid solution or an amino acid containing infusion regimen prior to administration. The rate of infusion depends on that of the carrier solution and should be adjusted according to the proportions of parenteral and enteral nutrition.

Duration of administration

The duration of use should not exceed 3 weeks.

Children

Safety and efficacy in children have not been established.

DIRECTION FOR USE

1. This product is a concentrated solution which should not be administered directly.
2. Perform the mixing of this concentrated solution with other amino acids solution under aseptic condition to maintain a good dispensing quality. Discard any remaining solution.
3. This product should be injected with a carrier solution. Mix one volume of this product with at least 5 volumes of compatible solution. (Ex: 100 ml of Kipo for IV Infusion + at least 500 ml of amino acids solution). Do not exceed the upper concentration limit of N(2)-L-alanine- L-glutamine, which is 3.5%.
4. Special attention before use:
 - a) Impact or vibration might occur during the transportation and caused breakage or cracks of containers, or loosened closures. Do not use the product if the above mentioned conditions are observed.
 - b) Do not use the product if the following conditions are observed: solution is no longer clear; changed color; foreign matters; other abnormal conditions.
5. Selection and insertion of infusion set:
 - a) Use the infusion set appropriately according to its instruction.
 - b) Use an infusion set with a filter. Insert the spike into the rubber stopper perpendicularly. Avoid twisting motion after insertion to prevent fragments caused by coring.
 - c) Visually inspect the solution and make sure there is no foreign matter prior to use.

WARNINGS AND PRECAUTIONS

Liver functions should be monitored routinely for patients with decompensated liver disease.

Blood electrolytes, osmotic pressure, water balance, acid-alkaline balance, and liver functions (alkaline phosphatase, ALT, AST), and potential hyperammonemia, should be controlled.

The levels of alkaline phosphatase, GOT, GPT, bilirubin, and acid-alkaline balance should be monitored.

Depending on the osmotic pressure of the mixed solution, this product is applied from central or peripheral veins. If the osmolarity is less than 800 mosmol/L, application from peripheral veins is acceptable. However, patient's age and condition, and the condition of peripheral veins should be considered.

The experience of using this product for more than 9 days is still limited.

Effects on ability to drive and use machines

Kipo for IV Infusion 200mg/mL is presumed to be safe or unlikely to produce an effect on the ability to drive or use machinery.

INTERACTIONS WITH OTHER DRUGS

No drug-drug interaction of Kipo for Infusion has been reported.

Incompatibility:

When mixing with the carrier solution, make sure the solution is injected in aseptic and completely mixed condition. Do not add any other medicinal product to this mixed solution. The mixed solution should be used immediately.

Storage of mixed solution is not allowed.

PREGNANT WOMEN AND NURSING MOTHERS

Application of this product to pregnant women and nursing mothers is not suggested due to lack of experiences and insufficient clinical data.

SIDE EFFECT

No known symptoms when correctly administered.

SYMPTOMS AND TREATMENT OF OVERDOSE

As with other infusion solutions, chills, nausea and vomiting can occur, when the infusion rate of Kipo for IV Infusion is exceeded. Infusion shall be stopped immediately in this case.

STORAGE

Store below 30°C. Protect from light and moisture. Do not freeze.

Keep out of reach of children.

Storage of mixed solution is not allowed.

Dispose the remaining product immediately after use. Once opened or inserted with a vent needle, the product should be used immediately. Do not use after being placed for a while.

SHELF-LIFE

Unopened - 24 months.

PACKAGING

The product is supplied in 50 mL and 100 mL glass bottles. Products less than 100 units are packaged in a box.

PRODUCT REGISTRATION HOLDER

Nang Kuang Pharm Sdn. Bhd.

A-07-3A Ekocheras, No. 693, Batu 5, Jalan Cheras,
56000 Kuala Lumpur Wilayah Persekutuan Kuala Lumpur, Malaysia

MANUFACTURED BY

Nang Kuang Pharmaceutical Co., Ltd.

No. 1001, 1001-1, Zhongshan Rd., Xinhua Dist., Tainan City 71243,
Taiwan (R.O.C.)

IMPORTED BY

Nano Medic Care Sdn. Bhd.

No. 1-6 & 1-12, 1st floor, Jalan Indah 2, Taman Indah, 84000 Muar,
Johor, Malaysia

REGISTRATION NO.:

MAL20026072ACZ

Revision Date: August 2024