

INTERACTION WITH OTHER MEDICAMENTS:

- Digoxin, cardiac glycosides
In patients under treatment with cardiac glycosides, care should be taken to keep the potassium concentration constant. In case of hyperkalaemia, the effect of cardiac glycosides may be weakened, and in case of hypokalaemia it may result in cardiac glycoside toxicity. Interactions might occur in the concurrent administration of other antiarrhythmics. Potassium administration must be very carefully discontinued in these patients.
- Drugs with the potential to induce hyperkalaemia
Care should be taken in the concurrent use of drugs containing potassium, potassium-sparing diuretics and drugs with a potential to induce hyperkalaemia such as Spironolactone, Triamterene, ACE inhibitors, Angiotensin II receptor antagonists, Non-steroidal anti-inflammatory agents, Ciclosporin, Tacrolimus and Suxamethonium. The concomitant administration of potassium-containing solutions and these drugs may lead to severe hyperkalaemia, which may in turn lead to cardiac arrhythmia. The concomitant use of sodium-retaining drugs (e.g. corticosteroids, nonsteroidal anti-inflammatory agents) may lead to oedema.
- Drugs leading to a decrease of the serum potassium level such as ACTH, corticosteroids and loop diuretics can increase the renal elimination of potassium.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

There are no or limited amount of data from the use of INFUSOL® K⁺0.3 NS in pregnant and lactating women. However, as all components of this medicinal product are naturally present in the body and their biochemical properties are well known, no toxic effects in relation to pregnancy and lactation are to be expected.

INFUSOL® K⁺0.3 NS can be used during pregnancy and breastfeeding if clinically needed.

For pregnancy, caution should be exercised in the presence of pre-eclampsia.

ADVERSE EFFECTS:

General disorder and administration site conditions

Not known: Local reactions at the infusion site, including local pain, venous irritation and occasionally thrombophlebitis may occur. In the case of undesirable effect(s), the infusion must be discontinued.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

INFUSOL® K⁺0.3 NS has no or negligible influence on the ability to drive and use machines.

OVERDOSE AND TREATMENT:

In case of overdose hyperkalaemia, hyperhydration, acid-base imbalances, oedema, particularly hypernatraemia, hyperchloraemia or potassium intoxication and electrolyte disorders can result. The symptoms of hyperkalaemia are primarily cardiovascular disorders and include hypotension, cardiac arrhythmia, heart block, ECG abnormalities with development of biphasic curves and cardiac arrest. Other symptoms include paresthesia of extremities, muscle or respiratory paralysis, areflexia, weakness and mental confusion. Rapid increase of the serum sodium level in patients with chronic hyponatraemia may lead to the osmotic demyelination syndrome.

Treatment:

Immediate interruption of the infusion, ECG monitoring, if necessary enhancement of urine flow and thus fluid and electrolyte excretion, administration of sodium bicarbonate and insulin. If insulin is given to increase cellular uptake of potassium, glucose should be given to avoid hypoglycemia. In patients with persistent ECG abnormalities e.g. calcium gluconate may be administered to antagonize the cardiotoxic effects of potassium. Hemodialysis or peritoneal dialysis may be required in patients with renal insufficiency.

INCOMPATIBILITIES:

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

CAUTION:

Single dose container. Discard all unused contents. Do not use if leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used.

STORAGE CONDITION: Do not store above 30°C.

SHELF LIFE:

2 years from manufacturing date in the proposed storage condition. Do not use after expiry.

DOSAGE FORM AND PACKAGING AVAILABLE:

500 mL x 20 LDPE bottle

MANUFACTURER / PRODUCT REGISTRATION HOLDER:



 **AIN MEDICARE SDN.BHD.**
Lot 4933, 4934 & PT2464, Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA.

HANDLING INSTRUCTION

AIN MEDICARE INTRAVENOUS INFUSION FLUID IN LOW DENSITY POLYETHYLENE (LDPE) BOTTLE





The bottle is made from injectable grade polyethylene.

PARTS OF THE PLASTIC BOTTLE



Top view after pull ring cap is removed. Please refer whichever applicable design.

INSTRUCTION FOR THE USAGE OF IV ADMINISTRATION SET & ADDITION OF DRUG



Date of Revision: 03.04.2019