

Lyso-B Lozenges

Lysozyme Hydrochloride 80mg, Pyridoxine Hydrochloride 40 mg

Product description: A white to off-white, round lozenges with menthol flavour, one side with scoreline and the other side with embossed logo AV.

Pharmacodynamics:

Lysozyme is an enzyme having a mucopolysaccharide lysing action. It enhances the effect of antibiotics to destroy or weaken cell bacteria. Lysozyme also accelerates immuno-bacteriolysis and activates the phagocytosis activity of leukocytes to destroy bacteria and viruses. Lysozyme is widely distributed in tears, nasal discharge, saliva and also breast milk. Lysozyme extracted from egg-white is used clinically for the intensification of natural defence mechanisms, restoration of damaged tissue and anti-inflammation.

The body needs pyridoxine hydrochloride in order to make cells of the immune system. Pyridoxine hydrochloride helps to improve immune response to the increase in production of antibodies and also helps in communicative interactions between cytokines and chemokines.

Pharmacokinetics:

In healthy men receiving a single 900 mg (potency) oral dose of lysozyme chloride, the peak plasma concentration occurred about 30 minutes to 1 hour after administration.

Pyridoxine hydrochloride is absorbed rapidly from the upper intestine regardless of the size of the dose given. Absorption may also occur from the ileum and to a small extent from the colon. There is a linear relationship between oral dose and the amount absorbed in normal animals and in those in which the distal small intestine has been resected. These observations suggest the possibility that pyridoxine may be absorbed by diffusion. Pyridoxine is rapidly converted in the liver to pyridoxine phosphate, pyridoxal phosphate and pyridoxamine phosphate via oxidation, this causes the release of pyridoxal and so pyridoxal phosphate to the general circulation where it reaches other organs chiefly as circulating pyridoxal. In urine, pyridoxine was excreted primarily unchanged with a small amount of only one metabolite, most likely 4-pyridoxic acid.

Indication: Adjuvant treatment of sore throat.

Recommended dose:

Adult: 1 lozenge to be sucked or dissolved in the mouth once or twice daily or according to doctor's prescription.

Contraindication:

Patients hypersensitive to egg or Vitamin B6.

Patients with a history of hypersensitivity to any ingredients of Lyso-B lozenges.

Warning and Precaution:

Patients with an allergic predisposition manifested as atopic dermatitis, bronchial asthma, drug allergy or food allergy to egg or pyridoxine hydrochloride (Vitamin B6). Abuse of pyridoxine may be responsible for sensory neuropathy that occurs with high doses of pyridoxine (> 200 mg/day) for a long period of time (months to years). This is usually reversible when the treatment is stopped. Please consult the doctor if condition does not improve after 5 days of treatment.

Pregnancy and Lactation: It is best not to use this medicine during pregnancy and breastfeeding.

Side effect:

May cause diarrhoea, anorexia, stomach discomfort, nausea and vomiting.

May cause hypersensitivity reactions such as skin rashes.

In rare case, shock may occur with administration of this drug. Therefore, patients should be carefully monitored during Lyso-B therapy.

Symptoms and Treatment of Overdose:

Symptoms: diarrhoea, anorexia, gastric discomfort.

Treatment: Stop administration.

Storage Condition: Store below 30°C.

Shelf-life: Three years from the date of manufacture.

Pack Size: 18x10's, 2x10's per box

Product registration holder: Advance Pharma Sdn Bhd

28, Jalan BP 6/6, Bandar Bukit Puchong, 47120 Puchong, Selangor.

Manufacturer: AV Manufacturing Sdn. Bhd (667760-V)

Lot 10621 (PT16700), Jalan Permata 2, Arab-Malaysian Industrial Park,
71800 Nilai, Negeri Sembilan, Malaysia.

Dec 2021, Ver. 02