

LIFE LINCO-SPEC PREMIX

FOR ANIMAL USE ONLY

Product Name :

LIFE LINCO-SPEC PREMIX

Name and Strength of Active Substance(s)

Each g contains Lincomycin 22 mg (equivalent to Lincomycin Hydrochloride Monohydrate 24.95 mg) and Spectinomycin 22 mg (equivalent to Spectinomycin Sulfate Tetrahydrate 33.26 mg).

Product Description

A white or off-white powder for use in animal feed.

Target Species

Pigs.

Pharmacodynamics

Lincomycin is a lincosamide antibiotic and is produced by *Streptococcus lincolnensis*. The mode of action of lincomycin is inhibition of protein synthesis at the ribosomal 50S sub-unit level. Lincomycin is largely used in pigs to control dysentery and mycoplasma infections.

Spectinomycin is an aminocyclitol antibiotic that is produced by *Streptomyces spectabilis*. The mode of action of spectinomycin is inhibition of protein synthesis via a 30S ribosomal target. It is a broad-spectrum drug with activity against gram-positive and some gram-negative bacteria, but little anaerobic activity. It has good activity against Mycoplasma.

Lincomycin and spectinomycin combination have been shown to be synergistic in vitro against mycoplasma.

Pharmacokinetics

Lincomycin is rapidly but incompletely absorbed when administered orally to animals. Lincomycin is well distributed in the body, with highest tissue concentration in the liver and kidneys, while very low levels are obtained in the CSF. Lincomycin can pass unchanged from the body via the bile and feces or urine or can be metabolized to the glucuronide, N-demethyl lincomycin, or icomycin sulfoxide forms by the liver.

Spectinomycin has a small volume of distribution. It is poorly absorbed following oral administration. Most of an oral dose is eliminated in the feces. Oral administration has been used for a local effect for treatment of diarrhea.

Target Species

Pigs

Indications

For the treatment and control of swine dysentery (Vibronic dysentery, bloody scours); salmonellosis (necrotic enteritis); enteritis associated with *E. coli* and as an aid in control of mycoplasmal pneumonia.

Route of Administration

To be mixed thoroughly in feed and administer orally.

Recommended Dosage

For the treatment and control of disease specified, complete feeds containing 44 grams of antibiotic activity per tonne should be fed.

Add 1 kg LIFE LINCO-SPEC PREMIX per tonne of complete feed will give 44g of antibiotic activity per tonne.

Direction of Use

LIFE LINCO-SPEC PREMIX must be thoroughly mixed in feeds before use. Medicated feed should constitute the sole ration of treated and growing swine.

Contraindications

None known.

Precautions

For use in pigs only. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feed containing Lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Lincomycin should not be used in animals known to be hypersensitive to the product. Do not use the water treatment and feed treatment simultaneously.

Drug Interactions

None known.

Pregnancy and Lactation

None known.

Adverse Effects

Excessive dosage may lead to an increased incidence of side effects which may include rectal prolapse. Occasionally, swine treated with Lincomycin, within the first two days after the onset of treatment, may develop diarrhoea and/or swelling of the anus. On the rare occasions, some pigs may show reddening of skin and irritable behavior. These conditions have been self-correcting within 5 to 8 days without discontinuing Lincomycin treatment.

Overdose and Treatment

If signs of overdose occur, discontinue treatment and, if necessary, treat symptomatically.

Withdrawal Period(s)

MEAT: DO NOT USE less than 1 day before slaughter for human consumption. Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withdrawal period.

Shelf-life

Shelf life as packaged for sale: 3 year

Shelf life after first opening : 24 hours

Shelf life after reconstitution : 24 hours

Maximum Residual Limit (MRL)

Lincomycin

Pig : Muscle 200 µg/kg, Liver 500 µg/kg, Kidney 1500 µg/kg, Fat 100 µg/kg, Skin with adhering fat 300 µg/kg [Ref. CODEX ALIMENTARIUS]

Spectinomycin

Pig : Muscle = 500 µg/kg, Fat = 2000 µg/kg, Liver = 2000 µg/kg, Kidney = 5000 µg/kg [Ref. CODEX ALIMENTARIUS]

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhi ubat daripada kanak-kanak.

Disposal of Containers

Emptied container must be safely disposed in trash according to local government regulations.

Packaging Available

25 kg in Low-density polyethylene plastic / polypropylene lined paper bag.

Manufacturer and Product Registration Holder

Life Biopharma Sdn. Bhd. (Company No. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2020-01-07