

DYNA CHARCOAL TABLET

MAL20032175XZ

DESCRIPTION:

TABLET

Colour : Black
Shape : Round and Biconvex
Coating : Colourless Coating

EACH TABLET CONTAINS:

Activated Charcoal.....250 mg

PHARMACODYNAMICS:

Adsorbent - Adsorbs the toxic substances ingested, thus inhibiting gastrointestinal absorption.
Antidiarrhoea - Adsorbs many toxic irritants that cause diarrhoea and gastrointestinal discomfort.
Antiflatulence - Adsorbs intestinal gas to relieve distension and discomfort.

PHARMACOKINETICS:

Absorption: Is not absorbed from the gastrointestinal tract.
Biotransformation: Is not metabolised.
Elimination: Through the intestinal tract.

INDICATIONS:

Treatment of flatulence and diarrhoea.

RECOMMENDED DOSE:

Adults : 2 - 4 tablets three to four times daily.
Children above 12 years : 1 - 2 tablets three to four times daily.
Drink plenty of fluid and maintain adequate salt intake.

ROUTE OF ADMINISTRATION: FOR ORAL USE ONLY

CONTRAINDICATIONS:

Dehydration - Rehydration therapy is necessary if signs of dehydration, such as dry mouth, excessive thirst, wrinkled skin, decreased urination, dizziness or light-headedness, are present; fluid loss may have serious consequences, such circulatory collapse and renal failure, especially in young children and the elderly.
Suspected parasite-associated diarrhoea - May make recognition of parasitic causes of diarrhoea more difficult; if parasitic agents are suspected pathogens, appropriate stool analysis should be performed prior to therapy with adsorbents.
Acute dysentery - sole treatment with adsorbent antidiarrhoeals may be inadequate; antibiotic therapy may be required.
Also contraindicated during the use of specific oral antidotes such as Methionine.

WARNINGS AND PRECAUTIONS:

Not recommended for treatment of diarrhoea in children under 6 years of age.

Elderly with diarrhoea: caution is recommended because of the risk of fluid and electrolyte loss.
Activated Charcoal may interfere with the absorption of other drugs, including antibiotics, when administered concurrently.
Appropriate fluid and electrolyte therapy should be given to protect against dehydration. Oral rehydration therapy which is the use of appropriate fluids including oral rehydration salts remains the most effective treatment for dehydration due to diarrhoea. The intake of as much of these fluids as possible is therefore imperative.

DRUG INTERACTIONS:

Activated Charcoal diminishes the action of lpecacuanha and other antiemetics when taken concomitantly by mouth. It also has the potential to reduce the absorption of many drugs from the gastrointestinal tract and simultaneous oral administration should therefore be avoided.

PREGNANCY AND LACTATION:

Pregnancy and breast-feeding: problems have not been documented.

SIDE EFFECTS:

Rarely, swelling or pain of abdomen.

PACKING/PACK SIZE(s):

Blister pack of 50x10's and 20x10's.

JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA

P11NT C002-05
Latest Revision: May 2021