

PRODUCT NAME

CAVUMOX 625 MG TABLET

NAME AND STRENGTH OF ACTIVE SUBSTANCE

Each tablet contains 500 mg amoxicillin (as amoxicillin trihydrate) and 125 mg clavulanic acid (as potassium clavulanate)

PRODUCT DESCRIPTION

White, oblong, biconvex film coated tablet

PHARMACODYNAMICS

CAVUMOX is a combination of amoxicillin (as trihydrate), a broad-spectrum antibiotic and clavulanic acid (as potassium salt), a progressive and irreversible inhibitor of β -Lactamase enzymes produced by many Gram-positive and Gram-negative bacteria.

The combination of amoxicillin/clavulanic acid extends the antibiotic spectrum of amoxicillin to include bacteria normally resistant to amoxicillin and β -lactam antibiotic.

CAVUMOX is bactericidal to a wide range of Gram-positive and Gram-negative bacteria includes β -lactamase penicillin resistant organisms.

PHARMACOKINETICS

The pharmacokinetics of the two components of CAVUMOX are closely matched. Peak serum levels of both occur about 1 hour after oral administration. Absorption of CAVUMOX is optimized at the start of a meal. Doubling the dosage of CAVUMOX approximately doubles the serum levels achieved. Both amoxicillin and clavulanate have low levels of serum binding; about 70% remains free in the serum.

INDICATION

CAVUMOX is an antibiotic agent with a notably broad spectrum of activity against the commonly occurring bacterial pathogens in general practice and hospital. The beta-lactamase inhibitory action of clavulanate extends the spectrum of amoxicillin to embrace a wider range of organisms, including many resistant to other beta-lactam antibiotics.

CAVUMOX should be used in accordance with local official antibiotic-prescribing guidelines and local susceptibility data.

CAVUMOX oral presentations for twice daily dosing, are indicated for short-term treatment of bacterial infections at the following sites:

- Upper respiratory tract infections, including ENT (e.g. tonsillitis, sinusitis, otitis media)
- Lower respiratory tract infections (e.g. acute exacerbation of chronic bronchitis, lobar and bronchopneumonia)
- Genito-urinary tract infections (e.g. cystitis, urethritis, pyelonephritis)
- Skin and soft tissue infections (e.g. boils, abscesses, cellulitis, wound infections)

- Bone and joint infections (e.g. osteomyelitis)
- Dental infections (e.g. dentoalveolar abscess)
- Other infections (e.g. septic abortion, puerperal sepsis, intra-abdominal sepsis)

Susceptibility to CAVUMOX will vary with geography and time. Local susceptibility data should be consulted where available, and microbiological sampling and susceptibility testing performed where necessary.

DOSAGE AND ADMINISTRATION

Dosage depends on the age, renal function of the patient and the severity of the infection.

To minimize potential gastrointestinal intolerance, administer at the start of a meal. The absorption of CAVUMOX is optimized when taken at the start of a meal.

Treatment should not be extended beyond 14 days without review.

Therapy can be started parenterally and continued with an oral preparation.

Tablets should be swallowed whole without chewing. If required, tablets may be broken in half and swallowed without chewing.

CAVUMOX tablets are not recommended in children of 12 years and under.

Adults and Children over 12 years

Mild to moderate infections: One CAVUMOX 625 mg tablet every 12 hours

Severe infections: One CAVUMOX 1 g tablet every 12 hours

Renal Impairment

No adjustment in dose is required in patient with creatinine clearance greater (CrCl) than 30 mL/min.

CAVUMOX 1 g tablet should only be used in patient with a CrCl rate of more than 30 mL/min.

CrCl 10-30 mL/min	One CAVUMOX 625 mg tablet every 12 hours
CrCl < 10 mL/min	One CAVUMOX 625 mg tablet every 24 hours
Haemodialysis	One CAVUMOX 625 mg tablet every 24 hours, plus a further one tablet during dialysis, to be repeated at the end of dialysis (as serum concentrations of both amoxicillin and clavulanic acid are decreased).

Hepatic Impairment

Dose with caution, monitor hepatic function at regular intervals.

CONTRAINDICATIONS

- CAVUMOX is contraindicated in patients with a history of hypersensitivity to beta-lactams (e.g. penicillins and cephalosporins).
- CAVUMOX is contraindicated in patients with a previous history of CAVUMOX-associated jaundice/hepatic dysfunction.

WARNINGS AND PRECAUTIONS

- Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients receiving penicillin (beta-lactam

antibiotic) therapy. Before initiating therapy with CAVUMOX, careful enquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, carbapenems or other beta-lactam agents. If an allergic reaction occurs, CAVUMOX must be discontinued immediately and appropriate alternative therapy instituted.

- CAVUMOX should be avoided if infectious mononucleosis is suspected since the occurrence of a morbilliform rash has been associated with this condition following the use of amoxicillin.
- Prolonged use may also occasionally result in overgrowth of non-susceptible organisms.
- Pseudomembranous colitis has been reported with the use of antibiotics and may range in severity from mild to life-threatening. Therefore, it is important to consider its diagnosis in patients who develop diarrhoea during or after antibiotic use. If prolonged or significant diarrhoea occurs or the patient experiences abdominal cramps, treatment should be discontinued immediately and the patient investigated further.
- Changes in liver function tests have been observed in some patients receiving CAVUMOX. The clinical significance of these changes is uncertain. CAVUMOX should be used with caution in patients with evidence of hepatic dysfunction.
- In patients with renal impairment, CAVUMOX dosage should be adjusted (See *Dosage and Administration*).
- In patients with reduced urine output, crystalluria has been observed very rarely, predominantly with parenteral therapy. During the administration of high doses of amoxicillin, it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria (See *Overdose and Treatment*).

INTERACTIONS

- Concomitant use of probenecid is not recommended. Probenecid decreases the renal tubular secretion of amoxicillin. Concomitant use with CAVUMOX may result in increased and prolonged blood levels of amoxicillin but not of clavulanate.
- Concomitant use of allopurinol during treatment with amoxicillin can increase the likelihood of allergic skin reactions. There are no data on the concomitant use of CAVUMOX and allopurinol.
- In common with other antibiotics, CAVUMOX may affect the gut flora, leading to lower oestrogen reabsorption and reduced efficacy of combined oral contraceptives.
- In patients receiving mycophenolate mofetil, reduction in pre-dose concentration of the active metabolite mycophenolic acid (MPA) of approximately 50% has been reported following commencement of oral amoxicillin plus clavulanic acid. The change in pre-dose level may not accurately represent changes in overall MPA exposure.

PREGNANCY AND LACTATION

Pregnancy

It was reported that prophylactic treatment with amoxicillin plus clavulanic acid may be associated with an increased risk of necrotising enterocolitis in neonates. As with all medicines, use should be

avoided in pregnancy, especially during the first trimester, unless considered essential by the physician.

Lactation

The medicinal product may be administered during the period of lactation. With the exception of the risk of sensitisation, associated with the excretion of trace quantities in breast milk, there are no detrimental effects for the infant.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Adverse effects on the ability to drive or operate machinery have not been observed.

UNDESIRABLE EFFECTS

Summary of Safety Profile

The adverse reactions presented below are classified by system organ and frequency (in decreasing order) from either clinical trial or post marketing experiences.

System Organ	Frequency	Adverse Reactions
Infections and infestations	Common	: Mucocutaneous candidiasis
Blood and lymphatic system disorders	Rare	: Reversible leucopenia (including neutropenia), thrombocytopenia
Immune system disorders	Very rare	: Angioneurotic oedema, anaphylaxis, hypersensitivity vasculitis
Nervous system disorders	Uncommon Very rare	: Dizziness, headache : Aseptic meningitis, convulsions (convulsions may occur in patients with impaired renal function or in those receiving high doses)
Gastrointestinal disorders (in adults)	Very common Common	: Diarrhoea : Nausea, vomiting
Gastrointestinal disorders (in children)	Common	: Diarrhoea, nausea, vomiting
Gastrointestinal disorders (in all populations) ¹	Uncommon Very rare	: Indigestion : Black hairy tongue
Hepatobiliary disorders	Uncommon	: A moderate rise in AST and/or ALT has been noted in patients treated with beta-lactam class antibiotics, but the significance of these findings is unknown
Skin and subcutaneous tissue disorders ²	Uncommon Rare Very rare	: Skin rash, pruritus, urticaria : Erythema multiforme : Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous exfoliative-dermatitis
Renal and urinary disorders	Very rare	: Interstitial nephritis

¹ Nausea is more often associated with higher oral dosages. If gastrointestinal reactions are evident, they may be reduced by taking CAVUMOX at the start of a meal.

² If any hypersensitivity dermatitis reaction occurs, treatment should be discontinued.

OVERDOSE AND TREATMENT

Gastrointestinal (GI) symptoms and disturbance of the fluid and electrolyte balances may be evident.

GI symptoms may be treated symptomatically with attention to the water electrolyte balance.

Amoxicillin crystalluria, in some cases leading to renal failure, has been observed.

CAVUMOX can be removed from the circulation by haemodialysis.

STORAGE CONDITIONS

Store in a tight container and in dry place below 30°C.

SHELF LIFE

2 years

DOSAGE FORMS AND PACKAGING AVAILABLE

5 blister strips x 3 film-coated tablets per unit box

DATE OF REVISION OF PI

Date of revision: 25/03/2022

Manufacturer: SIAM BHEASACH CO., LTD.

123 Soi Chokechairuammitr, Vibhavadi-Rangsit Road, Chomphon, Chatuchak, Bangkok 10900 and 9

Soi Chokechairuammitr 3, Vibhavadi-Rangsit Road, Dindang, Dindang, Bangkok 10400, Thailand.

Product Registration Holder, Importer: SPG PHARMA (MALAYSIA) SDN. BHD.

Suite 1701, Block A, Damansara Intan e-Business Park,

No. 1, Jalan SS20/27, 47400 Petaling Jaya, Selangor Darul Ehsan, Malaysia.